

Supercritical CO₂ extraction of fatty acids from cotton seeds (*Linum usitatissimum*): Experimental design and gas chromatography-mass

M. Sanayenaderi^{1,*}, M. Qomi¹, M. Salehi Sormaghi²

¹Department of Medicinal Chemistry (Pharmaceutical Sciences Research Center), Islamic Azad University of Pharmaceutical Sciences, Tehran, Iran

²Department of Pharmacognosy, Islamic Azad University of Pharmaceutical Sciences, Tehran, Iran

Background and Aims: In this present study, Supercritical fluid extraction (SFE) was developed for extraction of seeds oils from Cotton seeds. This method was used on a large scale for the production of essential oils and pharmaceutical products from plants.

Methods: The experimental parameters of SFE such as pressure, temperature, modifier volume, static and dynamic extraction time were optimized using a central composite design after a 2ⁿ-1 fractional factorial design. The chemical compositions of the SFE extract were identified by GC-MS. A SFE model suprexmts/225 and a GC-MASS 5890 hp Was used in this study.

Results: The major components of SFE extracted oils in optimum conditions were: Hexadecanoic acid, 8, 11-Octadecadienoic acid, Octadecanoic acid, Oleic acid, Eicosenoic acid, Extraction recovery based on the SFE varied in the range of 0.31–9.47% (w/w) under different conditions.

Conclusions: These results confirm supercritical fluid extraction compared with other extraction method is relatively rapid because of the low viscosities and high diffusivities associated with supercritical fluids. In this work, under the optimum conditions several fatty acids from cotton seeds with high recovery were extracted and identified by using SFE-GC-MASS.

Keywords: Supercritical fluid extraction (SFE); GC-MASS; Cotton seeds