

Protective effects of hydroalcoholic extract of *Stachys pilifera* on paracetamol-induced nephrotoxicity in female rats

Mohammad Reza Rabani¹, Nahid Azarmehr², Zahra Moslemi², Heibatollah Sadeghi¹, Hossein Amini-Khoei³, and Amir Hossein Doustimotlagh^{1,4,*}

¹Medicinal Plants Research Center, Yasuj University of Medical Sciences, Yasuj, I.R. Iran.

²Student Research Committee, Yasuj University of Medical Sciences, Yasuj, I.R. Iran.

³Medicinal Plants Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, I.R. Iran.

⁴Department of Clinical Biochemistry, Faculty of Medicine, Yasuj University of Medical Sciences, Yasuj, I.R. Iran.

Abstract

Background and purpose: *Stachys pilifera* is used in traditional medicine due to its antioxidant, anti-inflammatory, and antimicrobial effects. The goal of this study was to examine the renoprotective activity of the hydroalcoholic extract of aerial parts of *S. pilifera* on paracetamol (PCM)-induced nephrotoxicity.

Experimental approach: The Wistar female rats were randomly divided into four groups including control, PCM, *S. pilifera* hydroalcoholic extract (SPE), and PCM + SPE. The animals received SPE (500 mg/kg) for one week and PCM (3 g/kg) on the 6th day orally. Kidney function tests and oxidant/antioxidant markers were determined in serum and tissue homogenate, respectively. Protein and mRNA levels of TNF- α , as well as hematoxylin and eosin staining, were assessed in the kidney tissue.

Findings/Results: Treatment with SPE in the PCM group significantly decreased blood urea nitrogen and creatinine against the merely PCM rats ($P < 0.05$). The amount of nitric oxide metabolite and superoxide dismutase activity in the group receiving SPE showed a significant increase compared to PCM rats ($P < 0.05$). A significant difference in TNF- α levels between the groups was not observed. Histological changes were improved in the rats treated with SPE.

Conclusion and implications: Totally, our findings showed that SPE can inhibit PCM nephrotoxicity by enhancing kidney function markers, antioxidant status, and histological changes. Though, more researches are required to estimate the possible mechanism of SPE.

Keywords: Antioxidant; Nephrotoxicity; Paracetamol; *Stachys pilifera*.

INTRODUCTION

The kidneys are dynamic organs that are exposed to toxic damage due to their unique biochemical and physiological properties (1). Paracetamol (PCM, N-acetyl-para-aminophenol) or acetaminophen is a drug used as a painkiller and to control fever. It is safe in common doses, but overdose can induce hepatotoxicity and lead to nephrotoxicity if it progresses (2). However, acute kidney damage can also occur in the absence of liver damage and lead to death in humans and animals (3). Renal impairment is seen in 1 to 2% of patients receiving high-dose of PCM (4). The liver,

kidneys, and intestines are the most important organs involved in PCM metabolism (5). Following a therapeutic dose, PCM is inactivated mainly by conjugation with glucuronide and sulfate, and a small amount of it is oxidized to the N-acetyl-p-benzoquinone-imine (NAPQI) metabolite. This reactive metabolite is detoxified through glutathione (GSH), but in case of an overdose of PCM, the amount of NAPQI increases and causes depletion of GSH stores, oxidative stress, and eventually hepatorenal injury (6).

Access this article online



Website: <http://rps.mui.ac.ir>

DOI: 10.4103/1735-5362.327510

*Corresponding author: A.H. Doustimotlagh
Tel: +98-7433346070, Fax: +98-7433346071
Email: amirhossein.dousti@yums.ac.ir

In addition, PCM is converted to the nephrotoxic metabolite para-aminophenol by deacetylation in the kidneys, which causes necrosis of the renal cortex (4). Because oxidative stress is involved in the progress of kidney-liver damage caused by PCM, the use of natural compounds with antioxidant properties has been considered (5). Today, the renal protection effects of medicinal plants on PCM-induced nephrotoxicity have been proven (3).

The genus *Stachys* contains 300 species, which is one of the largest members of the Lamiaceae family and is distributed in tropical and subtropical regions (7). Thirty-four species of this genus are known in different regions of Iran, one of which is *Stachys pilifera*. Benth (8). In traditional medicine, several species of *Stachys* have been used as antinephritis, antidiarrhea, wound disinfectant, and anti-inflammation (9). In addition, the antibacterial, antioxidant, and cytotoxic properties of some species of *Stachys* have been proven (7). Flavonoids, iridoids, diterpenoids, and phenolic acids are secondary metabolites of various species of this genus (10). Also in our previous study, the antioxidant and hepatoprotective activity of the hydroalcoholic extract of this plant were observed in PCM-induced liver damage (11). The purpose of the current study was to evaluate the protective impact of *Stachys pilifera* hydroalcoholic extract (SPE) on PCM-induced nephrotoxicity in female rats.

MATERIAL AND METHODS

Preparation of the S. pilifera extract

S. pilifera plant was gathered in Yasuj, Iran, and identified by Dr. Jafari, from the Department of Botany, Yasuj University. A proof specimen (Voucher No. 1897) was deposited in the herbarium of the Department of Botany, Yasuj University, Kohgiluyeh, and Boyerahmad province, Yasuj, Iran. The dried aerial parts of *S. pilifera* were kept for 48 h at 25 °C in 70% ethanol. Then, the extract was filtered and the remnant was re-extracted with fresh ethanol (50%) for 24 h. The extract solution was concerted under condensed pressure at approximately 50 °C. The extraction yield of solid SPE was 20%.

Animals

This study used Wistar female rats that were approximately 200-250 g (adults, 8-week old). The rats were acquired from Shahrekord University of Medical Sciences, Shahrekord, Iran. The animals were maintained in a room under a controlled temperature (24 ± 2 °C) and were maintained in a 12/12-h light/dark cycle with free access to a normal diet and water to fed and drink *ad libitum*. All procedures used in the current study were carried out based on the "Principles of Laboratory Animal Care" (NIH Publication No. 86-23) and approved by the Ethics Committee of Yasuj University of Medical Sciences, Yasuj, Iran (Ethical code: IR.YUMS.REC.1398.012). All efforts were made to reduce animals from suffering and minimize the number of animals used in the study.

Experimental protocol

Twenty-four rats were randomly divided into four groups as follows: group I (the control): orally distilled water (vehicle) for one week; group II (PCM group): orally distilled water for one week and on the 6th day PCM (3 g/kg) (12,13); group III (SPE): orally ethanolic SPE (500 mg/kg) (8) for one week; and group IV (PCM + SPE): orally SPE (500 mg/kg) for one week and PCM (3 g/kg) on the 6th day. PCM was provided from Sigma Chemical Co (St Louis, MO, USA).

Specimen collection and biochemical assays

Twenty-four h after the administration of PCM, blood specimens were gathered from the cardiac puncture under ethyl ether anesthesia. They were centrifuged at 3500 g for 10 min and the serum was obtained for the assessment of biochemical parameters. After killing the rats, both kidneys were removed, washed entirely with ice-cold saline. One of them was kept in 10% formalin solution for histopathological analysis and the other was homogenized (10%, w/v) in phosphate-buffered saline (PBS; 10 mmol/l, pH 7.4) for biochemical analyses. Kidney homogenates were centrifuged at 10000 g for 5 min at 4 °C to determine nitric oxide (NO) metabolite, ferric reducing antioxidant power (FRAP), total thiols (T-SH), tumor necrosis factor alpha (TNF- α), and

antioxidant enzymes including GSH peroxidase (GPX), catalase (CAT), and superoxide dismutase (SOD).

Serum levels of urea and creatinine were evaluated using commercial kits (Pars Azmoon, Iran). The T-SH content was determined using the spectrophotometric method (14). Benzie and Strain procedure was applied to measure FRAP (14). Tissue NO metabolite (nitrite) was measured as indices of NO production according to the Griess reaction (15). Renal tissue homogenate was examined for the protein level of TNF- α according to the manufacturer's guidelines (ELISA kit, Kermania pars Gene, Kerman, Iran). For the determination of antioxidant enzymes activity (SOD, CAT, and GPX) in tissue homogenate, a routine kit (Zell Bio GmbH, Ulm, Germany) was used based on the manufacturer's procedure.

To explore mRNA TNF- α , the total RNA was extracted from kidney homogenate (RNX Plus kit, Sinaclon, Tehran, Iran) in line with the manufacturer's guidelines. First-strand complementary DNA (cDNA) was synthesized (cDNA Synthesis kit, Sinaclon, Tehran, Iran), and real-time polymerase chain reaction (RT-PCR) was done (Rotor Gene 3000 instrument, Bio-Rad, USA). The specific primer sequences were as follows: TNF- α and GAPDH (Table 1). PCR program was including the denaturation stage (95 °C for 15 S), annealing stage

(62 °C for 30 S), and elongation stage (72 °C for 30 S) in 40 cycles. The relative mRNA expression was calculated by the $2^{-\Delta\Delta Ct}$ formula.

Histological assessment

For light microscopic evaluation, kidney tissue portions were sectioned, fixed (in 10% formalin), dehydrated, and embedded. These samples were sliced into 5-mm thick segments, stained with hematoxylin & eosin, and observed under a light microscope by a pathologist who was blinded to the groups.

Statistical analysis

SPSS version 18 software was used to analyze the data. ANOVA followed by Tukey's tests were utilized to detect differences between groups. The values were reported as mean \pm SEM. *P* values < 0.05 were considered as a statistically noteworthy variation.

RESULTS

Biochemical markers

The results of this study demonstrated that serum urea and creatinine levels in the PCM group were considerably increased in comparison to the healthy rats. Treatment of PCM group with 500 mg/kg of SPE markedly decreased urea and creatinine levels compared to the merely PCM rats (Fig. 1A and B).

Table 1. The sequences of the primers for TNF- α and GAPDH.

| | Forward (5' to 3') | Reverse (5' to 3') |
|---------------|-----------------------|-----------------------|
| TNF- α | TGAGCACAGAAAGCATGATC | CATCTGCTGGTACCACCAGTT |
| GAPDH | AGTTCAACGGCACAGTCAAGG | AGACTCCACGACATACTCAGC |




Fig. 1. Impact of SPE (500 mg/kg) on (A) serum urea and creatinine (B) amounts in nephrotoxicity caused by PCM. Data are expressed as mean \pm SEM. **P* \leq 0.05, ***P* \leq 0.01, and ****P* \leq 0.001 indicate significant differences compared to the control group; #*P* \leq 0.05, ###*P* \leq 0.01, and ###*P* \leq 0.001 against the PCM group. PCM, Paracetamol, SPE, *Stachys pilifera* ethanolic extract.

TNF- α level

The renal tissue levels and mRNA expression of TNF- α slightly increased in the merely PCM rats against the control group but were not significant. Administration of SPE had no significant change on protein and mRNA levels of TNF- α in comparison to the PCM-treated animals (Fig. 2).

Oxidative stress parameters

As shown in Fig. 3, PCM-induced rats demonstrated a remarkable increment in FRAP level and a marked reduction in T-SH level in comparison to the control rats. In the PCM + SPE rats, FRAP and NO metabolite amounts were significantly increased, although, administration of SPE insignificantly increased T-SH content in

contrast to the PCM group.

Antioxidant enzymes evaluation

PCM significantly increased CAT activity in comparison to the control rats. Moreover, SOD activity was significantly augmented in the PCM + SPE group in contrast to PCM rats. However, the use of SPE had no considerable effect on the CAT and GPX enzymes activity. (Fig. 4).

Histological observation of kidney

Histological examination of renal tissue demonstrated that PCM causes cell necrosis and inflammation in renal glomeruli. White blood cells granulation was shown in the PCM rats. However, it was observed that the SPE could decrease these injuries (Fig. 5).




Fig. 2. The impact of SPE (500 mg/kg) on (A) the protein and (B) mRNA expression of TNF- α in nephrotoxicity caused by PCM. Data are expressed as mean \pm SEM. PCM, Paracetamol; SPE, *Stachys pilifera* ethanolic extract; TNF, tumor necrosis factor alpha.




Fig. 3. The effect of SPE (500 mg/kg) on oxidative stress parameters; (A) FRAP, (B) T-SH, (C) NO metabolite in nephrotoxicity caused by PCM. Data are expressed as mean \pm SEM. * $P \leq 0.05$ indicates significant differences compared to the control group; # $P \leq 0.05$ against the PCM group. PCM, Paracetamol, SPE, *Stachys pilifera* ethanolic extract; FRAP, ferric reducing antioxidant power; T-SH, total thiols; NO, nitric oxide.




Fig. 4. The impact of SPE (500 mg/kg) on the activity of the antioxidant enzymes of (A) CAT, (B) SOD, and (C) GPX in nephrotoxicity caused by PCM. Data are expressed as mean \pm SEM. * $P \leq 0.05$ indicates significant differences compared to the control group; # $P \leq 0.05$ against the PCM group. PCM, Paracetamol, SPE, *Stachys pilifera* ethanolic extract; CAT, catalase; SOD, superoxide dismutase; GPX, glutathione peroxidase.




Fig. 5. Histological examination of kidney tissues. (A) Control group; (B) paracetamol; (C); *Stachys pilifera* ethanolic extract; and (D) *Stachys pilifera* ethanolic extract + paracetamol.

DISCUSSION

PCM is one of the most effective analgesics and antipyretics belonging to the class of paraminophenol and nonsteroidal anti-inflammatory drugs (16). Although safe in therapeutic doses, overdose is the leading cause of liver damage among all drug toxicities (17). PCM-induced renal toxicity is mainly manifested after liver injury, but renal impairment in the absence of liver damage can also occur (1). Studies have shown that PCM-induced renal toxicity leads to acute kidney injury and death in experimental animals (18). Previous studies have demonstrated the renal-protection effects of medicinal plants on PCM-induced nephrotoxicity. In most of these studies, a dose of PCM in the range of 400 to 2000 mg/kg was used orally or intraperitoneally (3,19). Since plants belonging to the genus *Stachys* contain compounds such as flavonoids and phenolic acids, they can have an antioxidant role (9,10). Phenolic groups react with reactive oxygen species (ROS) as hydrogen donors and neutralize them (20).

Therefore, this study was performed to evaluate the impact of SPE on renal injury caused by PCM in rats.

Damage caused by drugs such as PCM in the kidney is characterized by tubular necrosis followed by crystalline nephropathy, uremia, glomerular hemodynamics, and inflammation (21). Increased serum levels of urea and creatinine are important diagnostic indicators for renal dysfunction because, in kidney damage, the rate of production of these substances increases relative to the rate of their clearance (22). The findings of the current study indicated that following the PCM administration (3 g/kg), serum urea and creatinine amounts increased significantly in comparison to the control group, which could indicate kidney tissue damage. These results were consistent with the findings of our previous study (13). Similar results were obtained in the previous studies by ingestion of 2 g/kg PCM (23). We observed that urea and creatinine levels were remarkably reduced in the group receiving SPE at a dose of 500 mg/kg. In accordance with the finding of the present

study, Sadeghi *et al.* showed that ethanolic extract of SPE at a dose of 500 mg/kg reduced urea and creatinine in cisplatin-induced renal damage (8). Therefore, it can be said that SPE is effective in preventing the progression of PCM-induced kidney damage.

As mentioned, the oxidative stress following the use of PCM can lead to severe kidney damage (3). Oxidative stress manifests itself as an imbalance between the production of oxidants, including ROS, and antioxidant compounds, leading to oxidative damage to proteins, lipids, nucleic acids, and ultimately destruction of tissue integrity (24). In our body, the antioxidant defense system traps ROS to inhibit oxidative stress (25). Among various antioxidant markers, SOD and CAT act as the chief enzymes in the elimination of ROS (25). In the present study, according to our previous investigation, consumption of PCM increased CAT enzyme activity (13). Although this change may be unexpected based on other studies (17,18), one study suggested that an increase in CAT could be due to the overproduction of free radicals (26). Thus, an augmentation in CAT in the present study is probably a compensatory response to an increase in free radical production after PCM. The use of SPE had no effect on CAT activity, although it significantly increased SOD activity. SOD is the main enzyme in reducing free radicals, including superoxide anions (11). Intracellular oxidative stress begins with the formation of superoxide, which is converted to oxygen and hydrogen peroxide by cytosolic or mitochondrial SOD (27). Semnani *et al.* showed that some *Stachys* species have antioxidant activity due to the reduction of superoxide anions (28). In a study performed by Sadeghi *et al.* it was observed that SPE with its scavenging properties has an antioxidant role against cisplatin-induced nephrotoxicity (8). Therefore, in the present study, increasing SOD activity in the SPE group may play a protective role against PCM-induced renal impairment through free radical scavenging activity and removal of superoxide anions.

T-SH groups of proteins are sensitive oxidative markers that are involved in the antioxidant defense system. T-SH contains protein thiol groups and GSH (29). Consistent

with our previous study (13), in this work, it was observed that in rats treated with PCM, T-SH levels were significantly reduced, which could be due to GSH depletion by NAPQI. Similar to our previous study (8,11) T-SH level enlarged in the SPE group in contrast to the PCM group. The FRAP method is a sensitive test for measuring the antioxidant power of biological fluids (30). In this study, it was observed that FRAP levels increased in both PCM and SPE-treated groups. Nitric oxide as an endogenous vasodilator is involved in the physiology and normal blood flow of the kidney and has been shown to reduce kidney damage in renal disease (31). In the present study, the amount of NO metabolite was slightly decreased in the PCM rats in contrast to the control group. Nagappan *et al.* observed that NO was reduced in indomethacin-induced renal injury *via* inhibiting NO synthase enzyme activity (32). Meng *et al.* showed that a decrease in NO in rats with urinary tract obstruction promotes the development of fibrosis in the renal tubules. However, they observed that consumption of *Astragalus membranaceus* and *Angelica sinensis* was able to increase NO levels by scavenging ROS and keeping NO (33). Therefore, in this study, it was suggested that the increase in NO metabolite following the injection of SPE is due to the trapping of free radicals by this plant.

Inflammation has been shown to be involved in the pathogenesis of PCM-induced renal damage (34). The nuclear factor kappa-B (NF- κ B) is a transcription factor that regulates immune responses and inflammatory diseases in many tissues (35). The NF- κ B pathway, which is involved in the activation of proinflammatory cytokines such as interleukine-6, TNF- α , and interleukine-1 β , is activated following oxidative stress and stimulates transcription of these cytokines (6). In this work, TNF- α levels were measured to evaluate the inflammatory process in the kidney. It was observed that PCM consumption insignificantly increased TNF- α . In agreement with this study, Ozatik *et al.* observed an increase in TNF- α in PCM-induced nephrotoxicity (4). We observed that *SP* extract had no significant effect on TNF- α levels.

CONCLUSION

Briefly, in this study, it was observed that the toxic dose of PCM can induce kidney damage. It has been speculated that the SPE may inhibit the development of nephrotoxicity due to its antioxidant properties as well as the improvement of biochemical and histological parameters. Nevertheless, more studies are needed in the future to show the exact mechanism of the impact of SPE on renal injury caused by PCM.

Acknowledgements

This study was financially supported by the Vice-Chancellor for Research of Yasuj University of Medical Sciences, Yasuj, I.R. Iran through Grant No. 960393).

Conflict of interest statement

All authors declared no conflict of interest in this study.

Authors' contribution

M.R. Rabani and H. Sadeghi designed the study, N. Azarmehr, Z. Moslemi, and H. Amini-Khoei performed the study, as well as A.H. Doustimotlagh analyzed the data and wrote the manuscript.

REFERENCES

1. Dokumacıoğlu E, Iskender H, Aktaş MS, Hanedan B, Dokumacıoğlu A, Mazlum Şen T, *et al.* The effect of sulforaphane on the levels of serum cystatin-c in acetaminophen-induced nephrotoxicity in rats. *Dicle Med J.* 2016;43(3):383-389. DOI: 10.5798/diclemedj.0921.2016.03.0701.
2. Li C, Liu J, Saavedra JE, Keefer LK, Waalkes MP. The nitric oxide donor, V-PYRRO/NO, protects against acetaminophen-induced nephrotoxicity in mice. *Toxicology.* 2003;189(3):173-180. DOI: 10.1016/s0300-483x(03)00129-x.
3. Chinnappan SM, George A, Thaggikuppe P, Choudhary Y, Choudhary VK, Ramani Y, *et al.* Nephroprotective effect of herbal extract *Eurycoma longifolia* on paracetamol-induced nephrotoxicity in rats. *Evid Based Complement Alternat Med.* 2019;2019:4916519,1-6. DOI: 10.1155/2019/4916519.
4. Ozatik FY, Teksen Y, Kadioglu E, Ozatik O, Bayat Z. Effects of hydrogen sulfide on acetaminophen-induced acute renal toxicity in rats. *Int Urol Nephrol.* 2019;51(4):745-754. DOI: 10.1007/s11255-018-2053-0.
5. Ozioko OM, Ozioko US, Mba CE, Atuadu V, Egwuatu IA, Okoro A. Curative effect of aqueous leaf extract of *Solanum macrocarpon* on acetaminophen induced nephrotoxicity on adult wistar rats. *World J Pharm Res.* 2020;9(6):158-172. DOI: 10.20959/wjpr20206-17650.
6. Eraky SM, El-Magd NFA. Omega-3 fatty acids protect against acetaminophen-induced hepatic and renal toxicity in rats through HO-1-Nrf2-BACH1 pathway. *Arch. Biochem Biophys.* 2020;687:108387. DOI: 10.1016/j.abb.2020.108387.
7. Kokhdan EP, Sadeghi H, Ghafoori H, Sadeghi H, Danaei N, Javadian H, *et al.* Cytotoxic effect of methanolic extract, alkaloid and terpenoid fractions of *Stachys pilifera* against HT-29 cell line. *Res Pharm Sci.* 2018;13(5):404-412. DOI: 10.4103/1735-5362.236833.
8. Sadeghi H, Mansourian M, Panahi kokhdan E, Salehpour Z, Sadati I, Abbaszadeh-Goudarzi K, *et al.* Antioxidant and protective effect of *Stachys pilifera Benth* against nephrotoxicity induced by cisplatin in rats. *J Food Biochem.* 2020;44(5):e13190. DOI: 10.1111/jfbc.13190.
9. Panahi Kokhdan E, Ahmadi K, Sadeghi H, Sadeghi H, Dadgary F, Danaei N, *et al.* Hepatoprotective effect of *Stachys pilifera* ethanol extract in carbon tetrachloride-induced hepatotoxicity in rats. *Pharm Biol.* 2017;55(1):1389-1393. DOI: 10.1080/13880209.2017.1302484.
10. Garjani A, Maleki N, Nazemieh H. Effects of hydroalcoholic extract from aerial parts of the sterile stems of *Stachys inflata* on myocardial infarct size in rats. *Iran J Pharm Sci.* 2004;3(3):165-170. DOI: 10.22037/IJPR.2010.595.
11. Mansourian M, Mirzaei A, Azarmehr N, Vakilpour H, Kokhdan EP, Doustimotlagh AH. Hepatoprotective and antioxidant activity of hydroalcoholic extract of *Stachys pilifera. Benth* on acetaminophen-induced liver toxicity in male rats. *Heliyon.* 2019;5(12):e03029,1-5. DOI: 10.1016/j.heliyon.2019.e03029.
12. Cristani M, Speciale A, Mancari F, Arcoraci T, Ferrari D, Fratantonio D, *et al.* Protective activity of an anthocyanin-rich extract from bilberries and blackcurrants on acute acetaminophen-induced hepatotoxicity in rats. *Nat Prod Res.* 2016;30(24):2845-2849. DOI: 10.1080/14786419.2016.1160235.
13. Ansari S, Azarmehr N, Barmoudeh Z, Moslemi Z, Ghahremani H, Mirzaei A, *et al.* Evaluation of the protective potential of hydroalcoholic extract of *Thymus daenensis* on acetaminophen-induced nephrotoxicity in rats. *Heliyon.* 2020;6(5):e03898. DOI: 10.1016/j.heliyon.2020.e03898.
14. Sadeghi A, Bastin AR, Ghahremani H, Doustimotlagh AH. The effects of rosmarinic acid on oxidative stress parameters and inflammatory cytokines in lipopolysaccharide-induced peripheral blood mononuclear cells. *Mol Biol Rep.* 2020;47(5):3557-3566. DOI: 10.1007/s11033-020-05447-x..

15. Doustimotlagh AH, Dehpour AR, Etemad-Moghadam S, Alaeddini M, Ostadhadi S, Golestani A. A study on OPG/RANK/RANKL axis in osteoporotic bile duct-ligated rats and the involvement of nitrenergic and opioidergic systems. *Res Pharm Sci.* 2018;13(3):239-249. DOI: 10.4103/1735-5362.228954.
16. Palani S, Kumar R, Kumar B. Effect of the ethanolic extract of *Indigofera barberi* (L.) in acute acetaminophen induced nephrotoxic rats. *New Biotechnol.* 2009;(25):S14. DOI: 10.1016/j.nbt.2009.06.989.
17. Ko JW, Shin JY, Kim JW, Park SH, Shin NR, Lee IC, et al. Protective effects of diallyl disulfide against acetaminophen-induced nephrotoxicity: a possible role of CYP2E1 and NF- κ B. *Food Chem Toxicol.* 2017;102:156-165. DOI: 10.1016/j.fct.2017.02.021.
18. Karaali HF, Fahmi RR, Borjac JM. Effect of *Ocimum basilicum* leaves extract on acetaminophen-induced nephrotoxicity in BALB/c mice. *J Complement Integr Med.* 2018;16(2):/j/jcim.2019.16.issue-2/jcim-2018-0111/jcim-2018-0111.xml. DOI: 10.1515/jcim-2018-0111.
19. Ezeonwu V, Dahiru D. Protective effect of bi-herbal formulation of *Ocimum gratissimum* and *Gongronema latifolium* aqueous leaf extracts on acetaminophen-induced hepato-nephrotoxicity in rats. *Am J Biochem.* 2013;3(1):18-23. DOI: 10.5923/j.ajb.20130301.03.
20. Kukić J, Petrović S, Niketić M. Antioxidant activity of four endemic *Stachys* taxa. *Biol Pharm Bull.* 2006;29(4):725-729. DOI: 10.1248/bpb.29.725.
21. Hussain Z, Khan JA, Arshad A, Asif P, Rashid H, Arshad MI. Protective effects of *Cinnamomum zeylanicum* L. (Darchini) in acetaminophen-induced oxidative stress, hepatotoxicity and nephrotoxicity in mouse model. *Biomed Pharmacother.* 2019;109:2285-2292. DOI: 10.1016/j.biopha.2018.11.123.
22. Reshi MS, Yadav D, Uthra C, Shrivastava S, Shukla S. Acetaminophen-induced renal toxicity: preventive effect of silver nanoparticles. *Toxicol Res.* 2020;9(4):406-412. DOI: 10.1093/toxres/tfaa040.
23. Haidara MA, Al-Ani B, Eid RA, Mohammed ME, Al-Hashem F, Dallak M. Acetaminophen induces alterations to the renal tubular ultrastructure in a rat model of acute nephrotoxicity protected by resveratrol and quercetin. *Int J Morphol.* 2020;38(3):585-591. DOI: 10.4067/S0717-95022020000300585.
24. Klimiuk A, Zalewska A, Sawicki R, Knapp M, Maciejczyk M. Salivary oxidative stress increases with the progression of chronic heart failure. *J Clin Med.* 2020;9(3):769-788. DOI: 10.3390/jcm9030769.
25. Ghosh J, Das J, Manna P, Sil PC. Acetaminophen induced renal injury via oxidative stress and TNF- α production: therapeutic potential of arjunolic acid. *Toxicology.* 2010;268(1-2):8-18. DOI: 10.1016/j.tox.2009.11.011.
26. Mansour MA, Nagi MN, El-Khatib AS, Al-Bekairi AM. Effects of thymoquinone on antioxidant enzyme activities, lipid peroxidation and DT-diaphorase in different tissues of mice: a possible mechanism of action. *Cell Biochem Funct.* 2002;20(2):143-151. DOI: 10.1002/cbf.968.
27. Jaeschke H, Ramachandran A. The role of oxidant stress in acetaminophen-induced liver injury. *Curr Opin Toxicol.* 2020;20-21:9-14. DOI: 10.1016/j.cotox.2020.03.003.
28. Morteza-Semnani K, Saeedi M, Shahani S. Antioxidant activity of the methanolic extracts of some species of *Phlomis* and *Stachys* on sunflower oil. *Afr J Biotechnol.* 2006;5(24):2428-2432.
29. Azarmehr N, Afshar P, Moradi M, Sadeghi H, Sadeghi H, Alipoor B, et al. Hepatoprotective and antioxidant activity of watercress extract on acetaminophen-induced hepatotoxicity in rats. *Heliyon.* 2019;5(7):e02072,1-5. DOI: 10.1016/j.heliyon.2019.e02072.
30. Gohari A, Hajimehdipoor H, Saeidnia S, Ajani Y, Hadjiakhoondi A. Antioxidant activity of some medicinal species using FRAP assay. *J Medicinal Plants.* 2011;10(37):54-60.
31. Huang A, Palmer LS, Hom D, Valderrama E, Trachtman H. The role of nitric oxide in obstructive nephropathy. *J Urol.* 2000;163(4): 1276-1281.
32. Nagappan AS, Varghese J, Pranesh GT, Jeyaseelan V, Jacob M. Indomethacin inhibits activation of endothelial nitric oxide synthase in the rat kidney: possible role of this effect in the pathogenesis of indomethacin-induced renal damage. *Chem Biol Interact.* 2014;221:77-87. DOI: 10.1016/j.cbi.2014.07.014.
33. Meng L, Qu L, Tang J, Cai SQ, Wang H, Li X. A combination of Chinese herbs, *Astragalus membranaceus* var. *mongholicus* and *Angelica sinensis*, enhanced nitric oxide production in obstructed rat kidney. *Vascul Pharmacol.* 2007;47(2-3):174-183. DOI: 10.1016/j.vph.2007.06.002.
34. Hua H, Ge X, Wu M, Zhu C, Chen L, Yang G, et al. Rotenone protects against acetaminophen-induced kidney injury by attenuating oxidative stress and inflammation. *Kidney Blood Press Res.* 2018;43(4):1297-1309. DOI: 10.1159/000492589.
35. Temel Y, Kucukler S, Yıldırım S, Caglayan C, Kandemir FM. Protective effect of chrysin on cyclophosphamide-induced hepatotoxicity and nephrotoxicity via the inhibition of oxidative stress, inflammation, and apoptosis. *Naunyn Schmiedebergs Arch Pharmacol.* 2020;393(3):325-337. DOI: 10.1007/s00210-019-01741-z.