



Anti-proliferative, anti-migration, and anti-invasion activity of novel hesperidin glycosides in non-small cell lung cancer A549 cells

Natwadee Poomipark, Titaporn Chaisin, and Jarunee Kaulpiboon*

Protein Research Laboratory, Division of Biochemistry, Department of Pre-Clinical Science, Faculty of Medicine, Thammasat University, Pathumthani 12120, Thailand.

Abstract

Background and purpose: Several attempts have been made to synthesize and investigate modified flavonoids to improve their potential anticancer efficacy. This study aimed to determine the *in vitro* anti-viability, anti-migration, and anti-invasive effects of two novel hesperidin glycosides, hesperidin glucoside (HG₁) and hesperidin maltoside (HG₂), compared to original hesperidin and diosmin.

Experimental approach: Inhibitory effects on normal (MRC5) and cancer (A549) cell viability of hesperidin glycosides were investigated by the trypan blue and MTS assays. A scratch assay determined the suppressive effects on cancer cell migration, and inhibition of cancer cell invasion was investigated through Matrigel™. The selectivity index (SI), a marker of cell toxicity, was also determined for A549 relative to MRC5 cells.

Findings/Results: The cell viability trypan blue and MTS assays showed similar results of the inhibition of A549 cancer cells; HG₁ and HG₂ had lower IC₅₀ than original hesperidin and diosmin. The SI of HG₁ and HG₂ was > 2 after 72-h culture. Investigation of cell migration showed that HG₁ and HG₂ inhibited the ability of gap closure in a time- and dose-dependent manner. The infiltration of the Matrigel™-coated filter by A549 cells was suppressed in the presence of HG₁ and HG₂. This result implied that HG₁ and HG₂ could inhibit cancer cell invasion.

Conclusion and implication: Our results suggest the inhibition of cancer cell migration and invasion in a time- and concentration-related manner with a favorable toxic profile. Moreover, HG₁ and HG₂ appeared potentially better agents than the original hesperidin for future anticancer development.

Keywords: Anticancer; Anti-invasion; Anti-migration; Anti-proliferation; Hesperidin glycosides.

INTRODUCTION

Cancer is one of the major public health problems in many countries around the world (1). Different environmental and genetic factors, such as oxidative stress, diet, radiation, smoking, etc. cause cancers in the human body. The meta-analyses showed a relationship between smoking with lung cancer risk, clearly seen for ever-smoking, current smoking, and even ex-smoking. It was stronger for squamous than adenocarcinoma and evident in both sexes (2). However, other factors such as asbestos, radon gas, air pollution exposure, and infections can participate in lung carcinogenesis (3). These factors play a crucial role in the pathophysiology of cancer (4).

There are several types of lung cancer therapies such as chemotherapy, radiation, surgery, and targeted therapy (3). The main treatment for early-stage disease is surgery which offers the best choice for long-term survival (5). Currently, various studies were established to develop new anticancer drugs. Among the new drugs, natural substances have been widely focused on (6). Flavonoids are naturally occurring phenolic compounds found in vegetables, plants, fruits, bark, and tea.

Access this article online



Website: <http://rps.mui.ac.ir>

DOI: 10.4103/1735-5362.383704

*Corresponding author: J. Kaulpiboon
Tel: +66-29269743, Fax: +66-29269711
Email: jkaulpiboon@gmail.com, kjarunee@tu.ac.th




Fig. 1. Chemical structures of (A) hesperidin, (B) diosmin, (C) hesperidin glucoside, and (D) hesperidin maltoside. Modified from Chaisin *et al.* (18).

They are thought to act as an anticancer, pro-apoptotic, and anti-proliferative effect in various cancer cell types (7) and some studies have sought to re-design and re-synthesize the flavonoid to increase some of these properties. A hydroxyl group substitution at the C-3 position on ring C and methylation substitution of free hydroxyls and 4-C=S are associated with the antiproliferative properties of flavonoids (8). Several flavonoids such as apigenin, anthocyanin, and quercetin have been reported to reduce cervical cancer cell viability and suppress cell metastasis and angiogenesis (9). Moreover, luteolin, one of the most prevalent flavonoids, was able to down-regulate the AKT signaling pathway and decrease the proliferation and migration of vascular smooth muscle cells (10).

Hesperidin is a natural flavanone glycoside. The hesperidin structure includes an aglycone unit, hesperetin, and a disaccharide, rutinose. Hesperidin has been demonstrated to suppress the viability of HeLa cells in a dose- and time-related manner and apoptosis in HeLa cells could be motivated by hesperidin *via* the acceleration of nuclear condensation and DNA fragmentation (11). In lung cancer studies, hesperidin induced apoptosis and suppressed the metastasis of cancer cells (12,13).

Our previous study on acceptor specificity (14) found that, among several flavonoids, hesperidin was the best acceptor for the

enzymatic synthesis of new flavonoid glycosides such as hesperidin glycosides (HG₁), hesperidin glucoside (HG₁) and hesperidin maltoside (HG₂) from p19bBC recombinant cyclodextrin glycosyltransferase (CGTase, E.C. 2.4.1.19). The basic properties and structures of both HG₁ and HG₂ were identified together with the related-structural compounds, hesperidin (Hes) and diosmin (Fig. 1) (15-18). So, the purpose of this work is to extend the knowledge of the HGs in the disease treatment of cancer by investigating the *in vitro* anti-proliferation, anti-migration, and anti-invasion properties of HG₁ and HG₂.

MATERIALS AND METHODS

Chemicals

Hesperidin and diosmin were purchased from Tokyo Chemical Industry Co., Ltd. (Tokyo, Japan). HG₁ and HG₂ were previously synthesized by cyclodextrin glycosyltransferase (CGTase, E.C. 2.4.1.19) and their molecular structures were determined (18). Dimethyl sulfoxide (DMSO) and cis-diamminedichloroplatinum II (cisplatin, DDP) were obtained from Sigma-Aldrich (USA). Fetal bovine serum (FBS) was from Himedia (India); Eagle's minimum essential medium (EMEM) and Ham's F-12 (Kaighn's modification) were purchased from Cassion (USA). Basement membrane matrix Matrigel®

was from Corning Life Sciences (USA) and cellTiter 96® AQueous one solution (MTS) was from Promega (USA). All other chemicals used were of analytical grade from Sigma-Aldrich (USA). Hesperidin, diosmin, and HGs were dissolved in DMSO.

Cell lines, culture conditions, and experimental groups

The MRC-5 (human lung fibroblast) and A549 (human lung carcinoma) cells were purchased from the American Type Culture Collection (ATCC, USA). The MRC-5 cells were cultured in EMEM supplemented with 1% penicillin/streptomycin and 10% FBS at 37 °C in 5% CO₂. The A549 cells were cultured in F-12K medium supplemented with 1% penicillin/streptomycin and 10% FBS at 37 °C in 5% CO₂.

In the present study, 6-main experimental groups were established as follows: control group containing 0.5% DMSO (the solvent of compounds); hesperidin groups (50, 100, and 150 µg/mL hesperidin); diosmin groups (50, 100, and 150 µg/mL diosmin), representing semi-synthetic hesperidin, and also diosmin has been reported to be a potential role in human diseases but only in a few lung cancer studies (19); HG₁ and HG₂ groups (50, 100, and 150 µg/mL). A chemotherapy medication DDP group (0.5, 1, and 2 µg/mL) was considered the positive control. The half maximal inhibitory concentration (IC₅₀) values of hesperidin (13) serve as a guide to decide the range of the concentration to be used in trypan blue and MTS assay for determining cell viability.

Determination of cell viability

Trypan blue assay

The MRC-5 and A549 cells were seeded at 5,000 cells/well on 96-well plates for 24 h and then treated with various treatments as mentioned before. The cells were cultured at 37 °C and 5% CO₂ for 24-72 h and cell growth was investigated at each time point. Cell viability was investigated by trypan blue dye exclusion assay. After trypsinization, quadruplicate wells of viable cells for each experimental group were counted on a hemocytometer. The growth curves were plotted, and the experiments were repeated at least three times. The concentration at which cell proliferation was inhibited by 50% (IC₅₀ value) was determined (GraphPad Prism

5.0, GraphPad Software, Inc., San Diego, CA, USA). In addition, the selectivity index (SI), indicating the safety of HGs for anticancer therapy was evaluated by obtaining the ratio of IC₅₀ for the non-cancer cell line to IC₅₀ for the cancer cell line (20).

MTS assay

The MRC-5 and A549 cells were seeded at 5,000 cells/well on 96-well plates for 24 h and then treated with various treatments as mentioned before. The cells were cultured at 37 °C and 5% CO₂ for 24, 48, and 72 h. After reaching each time point, MTS was added to each well. Then, the 96-well plates were incubated in a dark place at 37 °C and 5% CO₂ for 1 h and the absorbance was measured as the optical density at 490 nm using a microplate reader (Thermo Scientific, Multiskan GO, USA). The experiment was repeated three times. The cell viability was calculated using equation (1). Then, the IC₅₀ value and SI were calculated as described previously.

$$\text{Cell viability (\%)} = \frac{\text{Absorbance}_{\text{sample}}}{\text{Absorbance}_{\text{control}}} \times 100 \quad (1)$$

Determination of cell migration

The effects of hesperidin, diosmin, and HG₁ and HG₂ on cell migration were studied using the scratch assay. This method determines the movement of cells to close the gap between the scratch wound. The A549 cells were cultured in a 96-well plate to reach 90-100% confluence within 24 h. Then, a scratch was made on the cultured monolayer cells with a pipette tip and the scraped cells were cleaned with phosphate-buffered saline (PBS). The gap size of the wound was measured under the microscope (magnification ×400) as the width at the beginning of the scratch. Thereafter, the cells were further cultured in serum-free media containing 0-150 µg/mL of hesperidin, HGs, diosmin, and 1 µg/mL of DDP. The cells were incubated for 24-72 h at 37 °C in the 5% CO₂ incubator. The migration was observed using a phase-contrast microscope (magnification ×400) that also measured the width of the gaps at each time point. The experiment was repeated three times. The cell migration was determined by calculating % of wound closure from equation (2).

Gap closure (%) =

$$\frac{\text{Width at the beginning} - \text{width at a certain time}}{\text{Width at the beginning}} \times 100 \quad (2)$$

Determination of cell invasion

The Matrigel invasion assay was applied to determine the invasive capacity of the cells through Matrigel™, which acts as an extracellular matrix. Briefly, Matrigel™ was thawed, liquefied on ice, and then diluted with cold serum-free media. Then, Matrigel™ was added to a 96-Transwell® upper chamber and remained in a 37 °C incubator overnight to form a thin-layered gel. The A549 cells were suspended in serum-free media containing 0-150 µg/mL of hesperidin, diosmin, HG₁, and HG₂ and 1 µg/mL of DDP that was added to the Transwell® upper chamber. After that, 10% FBS (chemoattractant) was filled to the bottom of the lower chamber of the Transwell® plate. The cells were incubated at 37 °C in CO₂ for 24 and 48 h. After incubation, the media and remaining cells in the Transwell® upper chamber were cautiously removed and the Transwell® upper chamber was washed twice with PBS. The invasive cells attached to the Transwell® upper chamber were fixed on 3.7% paraformaldehyde and absolute methanol for 30 min, respectively. Then, the cells were stained with 50 µL of 1% crystal violet solution (Yd Diagnostics, Korea) for 15 min at room temperature and washed with PBS four times to eliminate the excess crystal violet dye. After that, the blue invasive cells on the Transwell® upper chamber were dried and counted under a microscope (magnification ×400) to enumerate the number of stained cells in five fields (21). The experiment was repeated three times. The cell invasion was determined by calculating % of invasion from equation (3).

$$\text{Invasion (\%)} = \frac{A_{\text{NU}} - A_{\text{NM}}}{A_{\text{NU}}} \times 100 \quad (3)$$

Where A_{NM} is the mean number of cells invading through the Matrigel™ matrix-coated permeable support membrane and A_{NU} is the mean number of cells migrating through the uncoated permeable support membrane.

Statistical analyses

Data were shown as mean ± SD from three independent experiments. Statistical analysis was accomplished using a one-way analysis of

variance (ANOVA) test followed by a post hoc Tukey test with the IBM SPSS Statistic version 26.0 (SPSS Corporation, Chicago, IL, USA). P -values ≤ 0.05 were considered significantly different.

RESULTS

The viability of MRC-5 and A549 cells

Inhibitory effects investigated by trypan blue exclusion assay

The cell survivability and suppression ratio in the 0.5% DMSO-treated cells were not markedly different from non-DMSO-treated cells (data were not shown). This implied that 0.5% DMSO, as the solvent of hesperidin, diosmin, HG₁, and HG₂, did not affect the viability of MRC-5 and A549 cells. Thus, the 0.5% DMSO-treated group was used as a control throughout the study. Hesperidin, diosmin, and HG₁ inhibited MRC-5 cells with an IC₅₀ value of > 150 µg/mL for 72 h, while HG₂ inhibited MRC-5 cells with an IC₅₀ value of 139.67 ± 3.18 µg/mL for 72 h. In contrast, the positive control (DDP) showed a good cell survival rate in MRC-5 cells at all concentrations. The IC₅₀ values at 72 h of hesperidin, diosmin, HG₁, HG₂, and DDP treatment of A549 cells were 92.90 ± 4.53, 97.66 ± 4.23, 88.85 ± 5.48, 87.35 ± 5.73, and 0.42 ± 0.04 µg/mL, respectively. It was evident that HG₁ and HG₂ had higher inhibitory activity than the original hesperidin. In addition, it was demonstrated that the cell proliferation following treatment with 150 µg/mL of hesperidin, diosmin, HG₁, and HG₂ at 24, 48, and 72 h was significantly reduced compared to the control in a time-dependent manner. In addition, the decreased cell proliferations were obtained with treatments at concentrations of 50, 100 and 150 µg/mL (Tables 1 and 2). So, it concluded that MRC-5 and A549 cell proliferation was reduced compared to the control cells in concentration-proportional manner. Furthermore, although the cell number increased with the treatments at different intervals, 24, 48, and 72 h, the MRC-5 and A549 cell proliferation rates were slowed down from time to time compared with those of the control. Moreover, the HG₂ treatment showed a selectivity index > 1.6, whereas the selectivity index of DDP at 72-h treatment was > 4. This implies that HGs have greater cytotoxic effects on normal cells compared to DDP.

Table 1. The effect of hesperidin, diosmin, HG₁, HG₂, and DDP on the number of MRC-5 cells. The data are expressed as the mean ± SD, n ≥ 3.

Sample	Concentration (µg/mL)	Number of cells			P period	P concentration
		24 h	48 h	72 h		
Control	0	14,166 ± 466	16,666 ± 365	22,916 ± 678		
Hesperidin	50	12,916 ± 498 ^a	14,583 ± 472 ^b	20,833 ± 404 ^{a,b}	$P_{c50, t24,48,72} \leq 0.001$	$P_{24} \leq 0.001$
	100	10,833 ± 473 ^{a,b}	13,333 ± 498	19,166 ± 523 ^{a,b}	$P_{c100, t24,48,72} \leq 0.001$	$P_{48} \leq 0.001$
	150	6,666 ± 662 ^{a,b}	11,250 ± 514 ^{a,b}	17,916 ± 508 ^{a,b}	$P_{c150, t24,48,72} \leq 0.001$	$P_{72} \leq 0.001$
Diosmin	50	11,666 ± 387 ^{a,b}	14,583 ± 489 ^b	19,583 ± 405 ^{a,b}	$P_{c50, t24,48,72} \leq 0.001$	$P_{24} \leq 0.001$
	100	9,583 ± 463 ^{a,b}	12,083 ± 723 ^a	17,500 ± 456 ^{a,b}	$P_{c100, t24,48,72} \leq 0.001$	$P_{48} \leq 0.001$
	150	6,666 ± 709 ^{a,b}	10,000 ± 630 ^{a,b}	15,833 ± 544 ^{a,b}	$P_{c150, t24,48,72} = 0.002$	$P_{72} \leq 0.001$
HG ₁	50	10,416 ± 598 ^{a,b}	13,333 ± 450	16,250 ± 602 ^{a,b}	$P_{c50, t24,48,72} \leq 0.001$	$P_{24} \leq 0.001$
	100	8,333 ± 523 ^{a,b}	10,000 ± 668 ^{a,b}	14,583 ± 582 ^b	$P_{c100, t24,48,72} \leq 0.001$	$P_{48} \leq 0.001$
	150	6,250 ± 846 ^{a,b}	8,333 ± 712 ^{a,b}	13,333 ± 608	$P_{c150, t24,48,72} \leq 0.001$	$P_{72} = 0.003$
HG ₂	50	9,583 ± 633 ^{a,b}	12,083 ± 604 ^a	15,000 ± 582 ^b	$P_{c50, t24,48,72} \leq 0.001$	$P_{24} \leq 0.001$
	100	7,500 ± 690 ^{a,b}	9,583 ± 757 ^{a,b}	13,333 ± 814	$P_{c100, t24,48,72} \leq 0.001$	$P_{48} \leq 0.001$
	150	5,000 ± 852 ^{a,b}	7,083 ± 809 ^{a,b}	10,833 ± 826 ^a	$P_{c150, t24,48,72} \leq 0.001$	$P_{72} \leq 0.001$
DDP	0.5	13,900 ± 302 ^b	15,666 ± 418 ^{a,b}	21,100 ± 488 ^{a,b}	$P_{c50, t24,48,72} \leq 0.001$	$P_{24} = 0.051$
	1	13,516 ± 414	15,400 ± 347 ^{a,b}	20,500 ± 465 ^{a,b}	$P_{c100, t24,48,72} \leq 0.001$	$P_{48} = 0.110$
	2	12,933 ± 396 ^a	14,933 ± 292 ^b	19,666 ± 389 ^{a,b}	$P_{c150, t24,48,72} \leq 0.001$	$P_{72} = 0.022$

HG, Hesperidin glucoside; DDP, diamminedichloroplatinum; ^a $P < 0.05$ indicates significant differences compared with the data of control after 24-h treatment; ^b $P < 0.05$ versus 50 µg/mL hesperidin after 24-h treatment.

Table 2. The effect of hesperidin, diosmin, HG₁, HG₂, and DDP on the number of A549 cells. The data are expressed as the mean ± SD, n ≥ 3.

Sample	Concentration (µg/mL)	Number of cells			P period	P concentration
		24 h	48 h	72 h		
Control	0	17,500 ± 465	22,500 ± 532	38,333 ± 616		
Hesperidin	50	15,000 ± 505 ^a	18,333 ± 522 ^b	30,416 ± 706 ^{a,b}	$P_{c50, t24,48,72} \leq 0.001$	$P_{24} \leq 0.001$
	100	10,833 ± 438 ^{a,b}	13,333 ± 630 ^{a,b}	17,500 ± 618 ^b	$P_{c100, t24,48,72} \leq 0.001$	$P_{48} \leq 0.001$
	150	7,500 ± 366 ^{a,b}	9,166 ± 414 ^{a,b}	12,083 ± 409 ^{a,b}	$P_{c150, t24,48,72} \leq 0.001$	$P_{72} \leq 0.001$
Diosmin	50	13,333 ± 444 ^{a,b}	18,333 ± 510 ^b	26,250 ± 392 ^{a,b}	$P_{c50, t24,48,72} \leq 0.001$	$P_{24} \leq 0.001$
	100	10,833 ± 487 ^{a,b}	13,750 ± 538 ^{a,b}	18,750 ± 426 ^{a,b}	$P_{c100, t24,48,72} \leq 0.001$	$P_{48} \leq 0.001$
	150	6,666 ± 539 ^{a,b}	8,750 ± 572 ^{a,b}	11,250 ± 559 ^{a,b}	$P_{c150, t24,48,72} \leq 0.001$	$P_{72} \leq 0.001$
HG ₁	50	13,333 ± 389 ^{a,b}	17,083 ± 394 ^b	27,500 ± 412 ^{a,b}	$P_{c50, t24,48,72} \leq 0.001$	$P_{24} \leq 0.001$
	100	10,833 ± 493 ^{a,b}	13,333 ± 468 ^{a,b}	17,083 ± 334 ^b	$P_{c100, t24,48,72} \leq 0.001$	$P_{48} \leq 0.001$
	150	6,250 ± 711 ^{a,b}	7,916 ± 678 ^{a,b}	10,416 ± 702 ^{a,b}	$P_{c150, t24,48,72} \leq 0.001$	$P_{72} \leq 0.001$
HG ₂	50	13,750 ± 495 ^{a,b}	16,666 ± 503 ^b	27,916 ± 659 ^{a,b}	$P_{c50, t24,48,72} \leq 0.001$	$P_{24} \leq 0.001$
	100	10,416 ± 589 ^{a,b}	11,666 ± 414 ^{a,b}	16,250 ± 712	$P_{c100, t24,48,72} \leq 0.001$	$P_{48} \leq 0.001$
	150	4,583 ± 792 ^{a,b}	6,250 ± 805 ^{a,b}	6,666 ± 780 ^{a,b}	$P_{c150, t24,48,72} \leq 0.001$	$P_{72} \leq 0.001$
DDP	0.5	14,000 ± 310 ^{a,b}	14,050 ± 523 ^{a,b}	16,666 ± 420 ^b	$P_{c50, t24,48,72} \leq 0.001$	$P_{24} \leq 0.001$
	1	11,250 ± 545 ^{a,b}	11,616 ± 590 ^{a,b}	12,833 ± 564 ^{a,b}	$P_{c100, t24,48,72} \leq 0.001$	$P_{48} \leq 0.001$
	2	8,583 ± 589 ^{a,b}	9,833 ± 650 ^{a,b}	8,983 ± 310 ^{a,b}	$P_{c150, t24,48,72} = 0.036$	$P_{72} \leq 0.001$

HG, Hesperidin glucoside; DDP, diamminedichloroplatinum; ^a $P < 0.05$ indicates significant differences compared with the data of control after 24-h treatment ^b $P < 0.05$ versus 50 µg/mL hesperidin after 24-h treatment.

Inhibitory effects investigated by MTS assay

The results at 72-h showed that hesperidin, diosmin, HG₁, and HG₂ have the potential to inhibit cell MRC-5 with IC₅₀ values of 137.58 ± 5.39, 149.69 ± 6.32, 149.44 ± 5.48, and 143.32 ± 5.53 µg/mL, respectively, compared to DDP of > 2 µg/mL. Expectedly, the DPP positive control showed a good cell survival rate in MRC-5 cells at every tested concentration and every time point of treatment. On the other hand, the IC₅₀ values of hesperidin, diosmin, HG₁, HG₂, and DDP

treatment of A549 cells were 58.66 ± 3.02, 106.31 ± 3.52, 54.57 ± 7.08, 49.44 ± 6.28, and 0.63 ± 0.03 µg/mL, respectively. It was evident that HG₁ had higher anti-proliferative properties than the original hesperidin, especially at 24- and 48-h treatment. This MTS result was in concordance with the trypan blue exclusion assay. Although the lowest inhibitory effect was changed from HG₂ in the trypan blue assay to HG₁ in the MTS assay, it was not a significant difference between HG₁ and HG₂ at every time point. Moreover, HG₂ treatment

showed a selectivity index of 2.90 toward this cell line relative to the MRC-5 cell line, whereas the selectivity index of DDP at 72-h was > 3. In addition, it was shown that the cell proliferation in the treatment of hesperidin, diosmin, HG₁, and HG₂ at 24, 48, and 72 h was a significant reduction in a concentration-related manner (Tables 3 and 4).

Inhibitory effects on migration of lung cancer cells

In the scratch assay, hesperidin, diosmin, HG₁, and HG₂ showed anti-migration activity

of A549 cells as shown in the relatively wider wound gaps than that of the control in Fig. 2A and B. The control group exhibited signs of cell migration resulting in a greater percentage of gap closure in control cells vs. the hesperidin-treated cells, as shown in Table 5. The highest percentage of gap closure at 24 h was shown in control at 13.23% while the lowest percentage of gap closure was shown in 150 µg/mL-treated HG₂ at 2.44% So, 24 h after scratching, the 150 µg/mL HG₂-treated cells showed slower migration than control cells.

Table 3. The effect of hesperidin, diosmin, HG₁, HG₂, and DDP on the viability of MRC-5 cells. The data are expressed as the mean ± SD, n ≥ 3.

Sample	Concentration (µg/mL)	Cell viability (%)			P period	P concentration
		24 h	48 h	72 h		
Control	0	100 ± 0.00	110.46 ± 1.21	103.29 ± 0.93		
	50	88.70 ± 5.30 ^a	85.83 ± 2.69 ^a	89.38 ± 6.40 ^{a,b}	$P_{c50,124,48,72} = 0.229$	$P_{24} \leq 0.001$
Hesperidin	100	54.82 ± 9.82 ^{a,b}	62.20 ± 4.84 ^{a,b}	67.65 ± 3.36 ^{a,b}	$P_{c100,124,48,72} = 0.081$	$P_{48} \leq 0.001$
	150	40.70 ± 8.00 ^{a,b}	43.36 ± 3.36 ^{a,b}	44.26 ± 6.40 ^{a,b}	$P_{c150,124,48,72} = 0.307$	$P_{72} \leq 0.001$
Diosmin	50	80.87 ± 8.46 ^a	77.99 ± 9.30 ^a	86.74 ± 1.38 ^a	$P_{c50,124,48,72} = 0.218$	$P_{24} = 0.002$
	100	54.73 ± 7.45 ^{a,b}	53.87 ± 9.19 ^{a,b}	70.97 ± 9.54 ^{a,b}	$P_{c100,124,48,72} = 0.004$	$P_{48} = 0.006$
HG ₁	50	75.64 ± 9.45 ^a	77.13 ± 1.74 ^{a,b}	72.09 ± 5.39 ^{a,b}	$P_{c50,124,48,72} = 0.365$	$P_{24} = 0.003$
	100	55.22 ± 7.69 ^{a,b}	55.44 ± 2.76 ^{a,b}	62.23 ± 9.78 ^{a,b}	$P_{c100,124,48,72} = 0.196$	$P_{48} = 0.001$
HG ₂	50	68.05 ± 9.20 ^{a,b}	68.39 ± 7.87 ^{a,b}	68.78 ± 4.63 ^{a,b}	$P_{c50,124,48,72} = 0.813$	$P_{24} = 0.013$
	100	50.98 ± 6.11 ^{a,b}	51.57 ± 3.03 ^{a,b}	63.19 ± 4.81 ^{a,b}	$P_{c100,124,48,72} = 0.016$	$P_{48} = 0.002$
DDP	0.5	98.57 ± 0.77 ^{a,b}	95.23 ± 1.52 ^a	93.06 ± 0.91 ^a	$P_{c50,124,48,72} = 0.007$	$P_{24} = 0.002$
	1	96.34 ± 0.81 ^a	92.63 ± 0.83 ^a	91.83 ± 1.22 ^a	$P_{c100,124,48,72} = 0.003$	$P_{48} = 0.005$
	2	93.76 ± 1.07 ^a	90.59 ± 0.47 ^a	89.79 ± 1.07 ^a	$P_{c150,124,48,72} = 0.009$	$P_{72} = 0.026$

HG, Hesperidin glucoside; DDP, diamminedichloroplatinum; ^aP < 0.05 indicates significant differences compared with the data of control after 24-h treatment; ^bP < 0.05 versus 50 µg/mL hesperidin after 24-h treatment.

Table 4. The effect of hesperidin, diosmin, HG₁, HG₂, and DDP on the viability of A549 cells. The data are expressed as the mean ± SD, n ≥ 3.

Sample	Concentration (µg/mL)	Cell viability (%)			P period	P concentration
		24 h	48 h	72 h		
Control	0	100 ± 0.00	124.87 ± 6.72	136.64 ± 8.40		
	50	86.42 ± 3.14 ^a	60.94 ± 5.53 ^{a,b}	53.35 ± 2.14 ^{a,b}	$P_{c50,124,48,72} = 0.024$	$P_{24} = 0.001$
Hesperidin	100	60.90 ± 7.39 ^{a,b}	52.10 ± 1.19 ^{a,b}	47.15 ± 3.07 ^{a,b}	$P_{c100,124,48,72} = 0.063$	$P_{48} = 0.017$
	150	49.88 ± 7.83 ^{a,b}	47.25 ± 4.31 ^{a,b}	37.60 ± 3.85 ^{a,b}	$P_{c150,124,48,72} = 0.245$	$P_{72} = 0.002$
Diosmin	50	72.32 ± 3.23 ^{a,b}	71.08 ± 1.37 ^{a,b}	54.89 ± 3.43 ^{a,b}	$P_{c50,124,48,72} = 0.025$	$P_{24} \leq 0.001$
	100	61.65 ± 4.07 ^{a,b}	57.62 ± 5.15 ^{a,b}	50.74 ± 1.19 ^{a,b}	$P_{c100,124,48,72} = 0.106$	$P_{48} = 0.003$
HG ₁	50	45.59 ± 4.59 ^{a,b}	47.32 ± 6.78 ^{a,b}	35.61 ± 5.95 ^{a,b}	$P_{c150,124,48,72} = 0.258$	$P_{72} = 0.003$
	100	63.86 ± 4.28 ^{a,b}	50.87 ± 8.91 ^{a,b}	52.13 ± 9.28 ^{a,b}	$P_{c50,124,48,72} = 0.249$	$P_{24} \leq 0.001$
HG ₂	50	47.46 ± 3.95 ^{a,b}	42.96 ± 5.73 ^{a,b}	44.11 ± 6.70 ^{a,b}	$P_{c100,124,48,72} = 0.101$	$P_{48} = 0.039$
	150	31.90 ± 2.34 ^{a,b}	31.22 ± 6.12 ^{a,b}	32.59 ± 5.27 ^{a,b}	$P_{c150,124,48,72} = 0.823$	$P_{72} = 0.044$
DDP	0.5	74.78 ± 6.22 ^{a,b}	57.74 ± 1.84 ^{a,b}	49.65 ± 5.07 ^{a,b}	$P_{c50,124,48,72} = 0.062$	$P_{24} = 0.005$
	100	58.58 ± 5.53 ^{a,b}	49.51 ± 3.97 ^{a,b}	37.79 ± 7.11 ^{a,b}	$P_{c100,124,48,72} = 0.095$	$P_{48} = 0.002$
DDP	0.5	49.02 ± 5.84 ^{a,b}	36.65 ± 5.80 ^{a,b}	32.19 ± 5.62 ^{a,b}	$P_{c150,124,48,72} = 0.150$	$P_{72} = 0.032$
	1	83.59 ± 2.30 ^a	72.06 ± 1.82 ^{a,b}	58.74 ± 2.28 ^{a,b}	$P_{c50,124,48,72} = 0.013$	$P_{24} \leq 0.001$
	2	60.93 ± 3.57 ^{a,b}	43.73 ± 2.78 ^{a,b}	32.21 ± 1.95 ^{a,b}	$P_{c100,124,48,72} = 0.014$	$P_{48} \leq 0.001$
	2	46.86 ± 2.31 ^{a,b}	30.31 ± 3.09 ^{a,b}	20.61 ± 1.64 ^{a,b}	$P_{c150,124,48,72} = 0.014$	$P_{72} \leq 0.001$

HG, Hesperidin glucoside; DDP, diamminedichloroplatinum; ^aP < 0.05 indicates significant differences compared with the data of control after 24-h treatment; ^bP < 0.05 versus 50 µg/mL hesperidin after 24-h treatment.




Fig.2. Representative images of the wounds at 0-72 h after treatment with 0 (control), 50, 100, and 150 µg/mL of (A) hesperidin and (B) hesperidin maltoside.

Table 5. The effect of hesperidin, diosmin, HG₁, HG₂, and DDP on the migration of A549 cells. The data are expressed as the mean ± SD, n ≥ 3.

Sample	Concentration (µg/mL)	Gap closure (%)				P period	P concentration
		0 h	24 h	48 h	72 h		
Control	0	0	13.23 ± 2.68	23.89 ± 2.86	33.24 ± 1.75		
Hesperidin	50	0	4.64 ± 0.58 ^a	8.81 ± 2.52 ^b	14.03 ± 0.83 ^b	$P_{c50, 24,48,72} = 0.016$	$P_{24} = 0.993$
	100	0	4.69 ± 0.63 ^a	6.76 ± 0.82 ^{a,b}	12.42 ± 0.48 ^{a,b}	$P_{c100, 24,48,72} \leq 0.001$	$P_{48} = 0.081$
	150	0	4.65 ± 0.45 ^a	5.16 ± 0.79 ^a	10.11 ± 1.64 ^b	$P_{c150, 24,48,72} = 0.013$	$P_{72} = 0.013$
Diosmin	50	0	4.88 ± 0.82 ^a	7.58 ± 1.07 ^{a,b}	10.82 ± 1.10 ^b	$P_{c50, 24,48,72} \leq 0.001$	$P_{24} = 0.026$
	100	0	3.97 ± 0.26 ^a	6.08 ± 0.87 ^a	8.24 ± 0.67 ^{a,b}	$P_{c100, 24,48,72} = 0.007$	$P_{48} = 0.050$
	150	0	3.32 ± 0.19 ^{a,b}	5.32 ± 0.64 ^a	6.84 ± 0.55 ^{a,b}	$P_{c150, 24,48,72} = 0.006$	$P_{72} = 0.003$
HG ₁	50	0	4.46 ± 0.73 ^a	7.30 ± 1.57 ^{a,b}	9.28 ± 1.49 ^{a,b}	$P_{c50, 24,48,72} = 0.012$	$P_{24} = 0.124$
	100	0	4.24 ± 0.78 ^a	5.25 ± 0.20 ^a	6.55 ± 0.30 ^{a,b}	$P_{c100, 24,48,72} = 0.023$	$P_{48} = 0.030$
	150	0	3.29 ± 0.12 ^{a,b}	4.37 ± 0.75 ^a	5.99 ± 0.26 ^{a,b}	$P_{c150, 24,48,72} = 0.019$	$P_{72} = 0.008$
HG ₂	50	0	2.98 ± 0.39 ^{a,b}	4.01 ± 0.41 ^a	5.37 ± 0.69 ^a	$P_{c50, 24,48,72} = 0.006$	$P_{24} = 0.055$
	100	0	3.07 ± 0.24 ^{a,b}	3.60 ± 0.14 ^{a,b}	4.03 ± 0.07 ^a	$P_{c100, 24,48,72} = 0.010$	$P_{48} = 0.011$
	150	0	2.44 ± 0.07 ^{a,b}	3.01 ± 0.17 ^{a,b}	3.71 ± 0.12 ^a	$P_{c150, 24,48,72} = 0.002$	$P_{72} = 0.005$
DDP	0.5	0	3.09 ± 0.25 ^{a,b}	4.36 ± 0.60 ^a	4.77 ± 0.19 ^a	$P_{c0.5, 24,48,72} = 0.021$	$P_{24} = 0.006$
	1	0	2.52 ± 0.16 ^{a,b}	3.18 ± 0.04 ^{a,b}	3.55 ± 0.42 ^a	$P_{c1.0, 24,48,72} = 0.043$	$P_{48} = 0.005$
	2	0	2.38 ± 0.06 ^{a,b}	2.92 ± 0.04 ^{a,b}	3.53 ± 0.16 ^{a,b}	$P_{c2.0, 24,48,72} = 0.004$	$P_{72} = 0.003$

HG, Hesperidin glucoside; DDP, diamminedichloroplatinum; ^a $P < 0.05$ indicates significant differences compared with the data of control after 24-h treatment; ^b $P < 0.05$ versus 50 µg/mL hesperidin after 24-h treatment.

Cell migration rate in the hesperidin, diosmin, HG₁, and HG₂ groups at 24, 48, and 72 h declined in a time-related manner as shown, especially HG₂ treatment, in the smaller relative change of % gap closure from time to time. The inhibition of cell migration also showed an increase in a concentration-dependent manner (Table 5). In addition, the anti-migration effect of HGs was substantially higher compared with both the control group and the original hesperidin. Moreover, the greatest effects on cell migration belonged to HG₂ at 150 µg/mL against the A549 cells, which was similar to that produced by DDP at 1.0 µg/mL.

Inhibitory effects on the invasion of A549 cells

The penetration of A549 cells through the

Matrigel™-coated filter was suppressed in the presence of hesperidin, diosmin, HG₁, and HG₂ (Fig. 3A and B show only the results of hesperidin and HG₂). The invasion percentage results of hesperidin, diosmin, HG₁, and HG₂ treatment at 150 µg/mL at 48 h were 33.89 ± 4.82, 27.60 ± 1.62, 24.40 ± 2.38, and 20.62 ± 2.35%, respectively, compared with 92.30 ± 2.25% invasion in the control. Furthermore, it was demonstrated that the cell invasion in the cells treated with hesperidin, diosmin, HG₁, and HG₂ at 24 and 48 h considerably declined in a time- and concentration-related manner (Table 6). Moreover, HG₁ and HG₂ showed a similar inhibition rate of invasion as the DDP positive control (Table 6).




Fig. 3. Transwell® assay was performed to determine A549 cell invasion. Images captured of representative invasive cells treated with (A) hesperidin, and (B) HG₂.

Table 6. Effect of hesperidin, diosmin, HG₁, and HG₂ on the invasion of A549 cells. The data are expressed as the mean \pm SD, $n \geq 3$.

Sample	Concentration ($\mu\text{g/mL}$)	Invasion (%)		P period	P concentration
		24 h	48 h		
Control	0	87.45 \pm 1.62	92.30 \pm 2.25		
Hesperidin	50	52.84 \pm 3.49 ^a	42.78 \pm 6.83 ^a	$P_{\text{c50, t24, 48}} = 0.233$ $P_{\text{c100, t24, 48}} = 0.001$ $P_{\text{c150, t24, 48}} = 0.115$	$P_{24} = 0.014$ $P_{48} = 0.264$
	100	47.11 \pm 5.60 ^a	37.06 \pm 6.26 ^{a,b}		
	150	39.03 \pm 1.50 ^{a,b}	33.89 \pm 4.82 ^{a,b}		
Diosmin	50	43.51 \pm 2.39 ^{a,b}	38.05 \pm 5.14 ^{a,b}	$P_{\text{c50, t24, 48}} = 0.075$ $P_{\text{c100, t24, 48}} = 0.002$ $P_{\text{c150, t24, 48}} = 0.024$	$P_{24} = 0.039$ $P_{48} = 0.028$
	100	38.60 \pm 3.31 ^{a,b}	31.39 \pm 2.71 ^{a,b}		
	150	34.79 \pm 3.59 ^{a,b}	27.60 \pm 1.62 ^{a,b}		
HG ₁	50	36.42 \pm 6.96 ^{a,b}	32.48 \pm 4.76 ^{a,b}	$P_{\text{c50, t24, 48}} = 0.090$ $P_{\text{c100, t24, 48}} = 0.698$ $P_{\text{c150, t24, 48}} = 0.063$	$P_{24} = 0.148$ $P_{48} = 0.135$
	100	31.55 \pm 5.70 ^{a,b}	28.82 \pm 4.84 ^{a,b}		
	150	26.04 \pm 3.13 ^{a,b}	24.40 \pm 2.38 ^{a,b}		
HG ₂	50	31.31 \pm 4.51 ^{a,b}	27.44 \pm 3.03 ^{a,b}	$P_{\text{c50, t24, 48}} = 0.045$ $P_{\text{c100, t24, 48}} = 0.448$ $P_{\text{c150, t24, 48}} = 0.011$	$P_{24} = 0.049$ $P_{48} = 0.039$
	100	27.54 \pm 2.97 ^{a,b}	24.90 \pm 1.91 ^{a,b}		
	150	22.64 \pm 1.97 ^{a,b}	20.62 \pm 2.35 ^{a,b}		
DDP	0.5	29.01 \pm 4.99 ^{a,b}	18.62 \pm 3.11 ^{a,b}	$P_{\text{c50, t24, 48}} = 0.011$ $P_{\text{c100, t24, 48}} = 0.008$ $P_{\text{c150, t24, 48}} = 0.012$	$P_{24} = 0.112$ $P_{48} = 0.205$
	1	26.12 \pm 4.22 ^{a,b}	16.13 \pm 2.66 ^{a,b}		
	2	20.91 \pm 2.05 ^{a,b}	14.42 \pm 1.57 ^{a,b}		

HG, Hesperidin glucoside; DDP, diamminedichloroplatinum; ^a $P < 0.05$ indicates significant differences compared with the data of control after 24-h treatment; ^b $P < 0.05$ versus 50 $\mu\text{g/mL}$ hesperidin after 24-h treatment.

DISCUSSION

The anticancer properties of the flavonoid glycoside in oranges (*Citrus sinensis* L.), hesperidin, and its flavone analog, diosmin, have exhibited anti-carcinogenic activities in

various studies (22). The anticancer effects of hesperidin are associated with its antioxidant and anti-inflammatory activities and its interactions with numerous cellular targets to suppress cancer cell proliferation by activating apoptosis and cell cycle arrest (23).

In the structure-activity relationship (SAR) of hesperidin, the ring B C-4' methyl replacement of hesperidin can motivate the ring B C-3' hydroxyl group, making hesperidin a better scavenger of free radicals (24). SAR for anticancer activity has interactions between the C2=C3 double bond (25). The substantial role of the C2=C3 double bond participates in molecular planarity and combination between rings C and A/B, which is crucial for powerful tumor suppression (26). Besides, adding glucose to the original structures of flavonoids or hesperidin, like our HG₁ and HG₂, increased their water solubility, bioavailability, and antioxidant activity (17) which could be a justification for why we found a greater anticancer activity of HG₁ and HG₂ compared to the original hesperidin regarding anti-viability, anti-migration, and anti-invasion properties. Although the anti-proliferation effect of HG was not as effective as the DDP positive control since the cytotoxic effect on normal cells of HGs was higher than DDP, its anti-proliferation properties would make a good promise. Moreover, the anti-migration and anti-invasion activities of HGs were comparable to those of DPP.

Cancer cell survival is suppressed by hesperidin over the mitochondrial apoptotic pathway and by inducing G0/G1 arrest in a time- and concentration-related manner. However, hesperidin does not have any adverse impacts on BEAS-2B normal cells (13). Xia *et al.* reported that the proliferation of A549 cancer cells was reduced by hesperidin, resulting in morphological alterations of apoptotic cells (13). They found that after treatment with various concentrations of hesperidin for 72 h, the A549 cell morphology changed and most of the cells treated with 1 µg/mL of DDP were apoptotic compared to that in the control group. Similarly, Cincin *et al.* found that hesperidin inhibits cell growth and motivates the programmed cell death pathway in two non-small cell lung cancer lines, A549 and NCI-H358, in a time- and concentration-related manner (7). They also demonstrated very low cytotoxicity of hesperidin in MRC-5 cells. Flavonoids have a dual action regarding reactive oxygen species homeostasis. They behave as antioxidants under normal cells and

are strong pro-oxidants in cancer cells activating programmed cell death pathways (27). Both antioxidant and pro-oxidant activities participate in flavonoid anticancer effects (27,28).

The communication of cancer cells with the extracellular matrix is crucial for metastasis, which is the primary reason for death in cancer patients. The repressive impact of hesperidin on migration and invasion of human non-small cell lung cancer cells may be mediated by the control of the chemokine stromal-cell derived factor-1, which is involved in promoting the neo-angiogenesis of cancer (12). In addition, hesperidin can suppress programmed death ligand 1, which is overexpressed in progressive cancer, and inhibit the activation of matrix metalloproteinases such as MMP-9 and MMP-2. These properties explain why hesperidin suppresses the metastatic phenotype and cell migration (29).

A flavonoid mixture tablet of hesperidin and diosmin (daflon) is marketed as a vasoprotective venotonic agent for the treatment of venous disease. This combination may prove useful as an anticancer agent and more work is needed on HG diosmin combinations to assess their potential anti-metastatic and anti-angiogenetic effects.

So far as we know, this study is the first to inform the suppressive effects of synthetic HG₁ and HG₂ on cancer cells. However, in-depth research is necessary to elucidate the underlying mechanisms of cancer as well as on the effects of HGs on cancer behavior in a physiologic environment to provide information for innovative-drug development in the future.

CONCLUSION

Our results suggest that new the HGs, HG₁ and HG₂, have more potential to inhibit cancer cells than the original hesperidin. They were effective against A549 cell lines and had a favorable SI score of > 2.0 relative to MRC-5 normal cells, suggesting a good toxicity profile. The suppression of cell viability, cell migration, and cell invasion by HG₁ and HG₂ was time- and concentration-dependent. Taken together, our new HGs have the potential as a new

alternative anticancer agent or may be used as a combination regimen, especially against metastases. More preclinical work is needed to ascertain whether HG₁ and HG₂ should be tested in humans.

Acknowledgments

This study was supported by Thammasat University Research Fund, Contract No. TUFT 63/2565. We also thank the Drug Discovery and Development Center, Center of Scientific Equipment for Advanced Research, Thammasat University, and the Research Group in Medical Biomolecules, Faculty of Medicine, Thammasat University.

Conflict of interest statement

The authors declared no conflict of interest in this study.

Authors' contributions

N. Poomipark and J. Kaulpiboon conceptualized the study. T. Chaisin and J. Kaulpiboon conducted the experiments. N. Poomipark and J. Kaulpiboon contributed to the methodology. T. Chaisin prepared the raw data file for analysis. N. Poomipark and J. Kaulpiboon analyzed the data and helped in the manuscript writing. The final version of the manuscript was approved by all authors.

REFERENCES

- Ma X, Yu H. Cancer issue: global burden of cancer. *Yale J Biol Med.* 2006;79(3-4):85-94. PMID: 17940618.
- Lee PN, Forey BA, Coombs KJ. Systematic review with meta-analysis of the epidemiological evidence in the 1900s relating smoking to lung cancer. *BMC Cancer.* 2012;12:385,1-90. DOI: 10.1186/1471-2407-12-385.
- Lemjabbar-Alaoui H, Hassan OU, Yang YW, Buchanan P. Lung cancer: biology and treatment options. *Biochim Biophys Acta.* 2015;1856(2):189-210. DOI: 10.1016/j.bbcan.2015.08.002.
- Devi KP, Rajavel T, Nabavi SF, Setzer WN, Ahmadi A, Mansouri K, *et al.* Hesperidin: a promising anticancer agent from nature. *Ind Crops Prod.* 2015;76:582-589. DOI: 10.1016/j.indcrop.2015.07.051.
- Sangha R, Price J, Butts CA. Adjuvant therapy in non-small cell lung cancer: current and future directions. *Oncologist.* 2010;15(8):862-872. DOI: 10.1634/theoncologist.2009-0186.
- Rayan A, Raiyn J, Falah M. Nature is the best source of anticancer drugs: indexing natural products for their anticancer bioactivity. *PloS One.* 2017;12(11):e0187925,1-12. DOI: 10.1371/journal.pone.0187925.
- Cincin ZB, Unlu M, Kiran B, Bireller ES, Baran Y, Cakmakoglu B. Anti-proliferative, apoptotic and signal transduction effects of hesperidin in non-small cell lung cancer cells. *Cell Oncol (Dordr).* 2015;38(3):195-204. DOI: 10.1007/s13402-015-0222-z.
- Ravishankar D, Watson KA, Greco F, Osborn HMI. Novel synthesised flavone derivatives provide significant insight into the structural features required for enhanced anti-proliferative activity. *RSC Adv.* 2016;6(69):64544-64556. DOI: 10.1039/C6RA11041J.
- Chen P, Xu R, Wang J, Wu Z, Yan L, Zhao W, *et al.* Starch biotransformation into isomaltooligosaccharides using thermostable alpha-glucosidase from *Geobacillus stearothermophilus*. *Peer J.* 2018;6:e5086,1-22. DOI: 10.7717/peerj.5086.
- Luo Y, Shang P, Li D. Luteolin: a flavonoid that has multiple cardio-protective effects and its molecular mechanisms. *Front Pharmacol.* 2017;8:692,. DOI: 10.3389/fphar.2017.00692,1-10.
- Wang Y, Yu H, Zhang J, Gao J, Ge X, Lou G. Hesperidin inhibits HeLa cell proliferation through apoptosis mediated by endoplasmic reticulum stress pathways and cell cycle arrest. *BMC Cancer.* 2015;15:682,1-11. DOI: 10.1186/s12885-015-1706-y.
- Xia R, Xu G, Huang Y, Sheng X, Xu X, Lu H. Hesperidin suppresses the migration and invasion of non-small cell lung cancer cells by inhibiting the SDF-1/CXCR-4 pathway. *Life Sci.* 2018;201:111-120. DOI: 10.1016/j.lfs.2018.03.046.
- Xia R, Sheng X, Xu X, Yu C. Hesperidin induces apoptosis and G0/G1 arrest in human non-small cell lung cancer A549 cells. *Int J Mol Med.* 2018;41(1):464-472. DOI: 10.3892/ijmm.2017.3250.
- Chaisin T, Kaulpiboon J, Poomipark N. Acceptor specificity of recombinant cyclodextrin glycosyltransferase from *Bacillus circulans* A11. *Sci Tech Asia.* 2018;23:86-97. DOI: 10.14456/scitechasia.2018.17.
- Kometani T, Terada Y, Nishimura T, Takii H, Okada S. Transglycosylation to hesperidin by cyclodextrin glucanotransferase from an alkalophilic *Bacillus* species in alkaline pH and properties of hesperidin glycosides. *Biosci Biotechnol Biochem.* 1994;58(11):1990-1994. DOI: 10.1271/bbb.58.1990.
- Kometani T, Nishimura T, Nakae T, Takii H, Okada S. Synthesis of neohesperidin glycosides and naringin glycosides by cyclodextrin glucanotransferase from an alkalophilic *Bacillus* species. *Biosci Biotechnol Biochem.* 1996;60(4):645-649. DOI: 10.1271/bbb.60.645.

17. Poomipark N, Chaisin T, Kaulpiboon J. Synthesis and evaluation of antioxidant and β -glucuronidase inhibitory activity of hesperidin glycosides. *Agr Nat Resour.* 2020;54(2):165-172. DOI: 10.34044/j.anres.2020.54.2.07.
18. Chaisin T, Rudeekulthamrong P, Kaulpiboon J. Enzymatic synthesis, structural analysis, and evaluation of antibacterial activity and α -glucosidase inhibition of hesperidin glycosides. *Catalysts.* 2021;11(5):532,1-17. DOI: 10.3390/catal11050532.
19. Huwait E, Mobashir M. Potential and therapeutic roles of diosmin in human diseases. *Biomedicines.* 2022;10(5):1076,1-25. DOI: 10.3390/biomedicines10051076.
20. Prayong P, Barusru S, Weerapreeyakul N. Cytotoxic activity screening of some indigenous Thai plants. *Fitoterapia.* 2008;79(7-8):598-601. DOI: 10.1016/j.fitote.2008.06.007.
21. Justus CR, Leffler N, Ruiz-Echevarria M, Yang LV. *In vitro* cell migration and invasion assays. *J Vis Exp.* 2014(88):51046,1-8. DOI: 10.3791/51046.
22. Manthey JA, Guthrie N. Antiproliferative activities of citrus flavonoids against six human cancer cell lines. *J Agric Food Chem.* 2002;50(21):5837-5843. DOI: 10.1021/jf020121d.
23. Aggarwal V, Tuli HS, Thakral F, Singhal P, Aggarwal D, Srivastava S, et al. Molecular mechanisms of action of hesperidin in cancer: recent trends and advancements. *Exp Biol Med (Maywood).* 2020;245(5):486-497. DOI: 10.1177/1535370220903671.
24. van Acker SA, van den Berg DJ, Tromp MN, Griffioen DH, van Bennekom WP, van der Vijgh WJ, et al. Structural aspects of antioxidant activity of flavonoids. *Free Radic Biol Med.* 1996;20(3):331-342. DOI: 10.1016/0891-5849(95)02047-0.
25. Murthy KNC, Kim J, Vikram A, Patil BS. Differential inhibition of human colon cancer cells by structurally similar flavonoids of citrus. *Food Chem.* 2012;132(1):27-34. DOI: 10.1016/j.foodchem.2011.10.014.
26. Krych-Madej J, Gebicka L. Catalase is inhibited by flavonoids. *Int J Biol Macromol.* 2013; 58:148-153. DOI: 10.1016/j.ijbiomac.2013.03.070.
27. Kopustinskiene DM, Jakstas V, Savickas A, Bernatoniene J. Flavonoids as anticancer agents. *Nutrients.* 2020;12(2):457,1-24. DOI: 10.3390/nu12020457.
28. Simunkova M, Barbierikova Z, Jomova K, Hudecova L, Lauro P, Alwasel SH, Alhazza I, Rhodes CJ, Valko M. Antioxidant vs. prooxidant properties of the flavonoid, kaempferol, in the presence of Cu (II) ions: a ROS-scavenging activity, Fenton reaction and DNA damage study. *Int J Mol Sci.* 2021;5;22(4):1619. DOI: 10.3390/ijms22041619.
29. Kongtawelert P, Wudtiwai B, Shwe TH, Pothacharoen P, Phitak T. Inhibitory effect of hesperidin on the expression of programmed death ligand (PD-L1) in breast cancer. *Molecules.* 2020;25(2):252,1-11. DOI: 10.3390/molecules25020252.