

Antibacterial, antifungal and cytotoxic evaluation of some new 2,3-disubstituted 4(3H)-quinazolinone derivatives

G.A. Khodarahmi^{1,2}, M. Rahmani Khajouei¹, G.H. Hakimelahi^{1,2,3}, D. Abedi⁴, E Jafari¹
and F. Hassanzadeh^{1,2,*}

¹Department of Medicinal Chemistry, School of Pharmacy and Pharmaceutical Science, Isfahan University of Medical Sciences, Isfahan, I.R.Iran.

²Isfahan Pharmaceutical Sciences Research Center, Isfahan, I.R.Iran.

³Institute of Chemistry, Academia Sinica, Nankang, Taipei, Taiwan.

⁴Department of Biotechnology and Isfahan Pharmaceutical Sciences Research Center, School of Pharmacy and Pharmaceutical Science, Isfahan University of Medical Sciences, Isfahan, I.R.Iran.

Abstract

In this study antibacterial, antifungal and cytotoxic effects of some new 2,3-disubstituted 4(3H)-quinazolinone derivatives have been evaluated. The *in vitro* antibacterial and antifungal tests of new synthesized compounds were performed using MABA method against six strains of bacteria (three Gram-positive and three Gram-negative) and three strains of fungi. Also Minimum Bactericidal Concentration (MBC) and Minimum Fungicidal Concentration (MFC) tests were performed. All synthesized compounds indicated mild antibacterial effects especially against Gram-negative bacteria. The most sensitive bacterium was *E. coli*. All strains of tested fungi were sensitive to the synthesized compounds mostly at 32 µg/ml and there was no significant differences in the sensitivity of the tested compounds. MBC and MFC data indicated that tested compounds act as bacteriostatic and fungistatic agents. Cytotoxic activity of the compounds was screened at 1, 10 and 100 µM concentrations against HeLa cells using the MTT colorimetric assay. While the synthesized compounds did not show significant cytotoxic activities, compounds **7a₃** and **7a₄** reduced cell viability to about 50% at 100 µM concentration. The present study revealed that most of the new synthesized compounds possess good antifungal effects and they could be considered as valuable candidates for further structural modification to design more potent antifungal agents.

Keywords: 4(3H)-Quinazolinone; Antibacterial; Antifungal; Cytotoxicity

INTRODUCTION

Currently one of the most important health problems is drug resistance due to the increased use and misuse of antibiotics and also lack of immunity in cases such as Acquired Immune Deficiency Syndrome (AIDS) and organ transplantation. Therefore, developing of new antimicrobial agents is unavoidable and many researchers have focused their efforts to find new and effective antibiotics from natural and synthetic sources. In addition, cancer is the second major life threatening disease in both developing and undeveloped countries. Although many advances have been made in the treatment of some types of cancers, the continued efforts of

the researchers to discover a new anticancer agent is also critically important (1).

Quinazolines are a group of fused heterocyclic compounds which have attracted interest of researchers due to their valuable biological properties including antifungal (2-4), antibacterial (4-6), and cytotoxic (7) activities.

There are many reports indicating that 2,3-disubstituted 4(3H)-quinazolinones have valuable antimicrobial properties. Structure activity relationship (SAR) studies have shown that positions 2 and 6 of quinazolinone structure are important and position 3 should be presented for better antimicrobial activities. The 3-substitution includes phenyl rings, bridged phenyl rings, heterocyclic rings and different aliphatic systems (8,9). Quinolones

*Corresponding author: F. Hassanzadeh, this paper is extracted from the Ph.D thesis No. 388303
Tel. 0098 311 7922575, Fax. 0098 311 6680011
Email: Hassanzadeh@pharm.mui.ac.ir

perform their antibacterial activities by inhibition of bacterial DNA gyrase which is an enzyme essential for bacterial DNA replication (10,11). Due to the structure similarity between quinazolinones and quinolones, antibacterial effects of quinazolinones could be a result of inhibitory activity against DNA replication. Also antibacterial activity was observed in a group of quinazoline derivatives as a result of DNA polymerase III inhibitory effect. The inhibition of chromosomal DNA replication results in bacterial cell death (12).

It is also reported that quinazoline derivatives bearing aromatic rings and some nitrogen containing groups in their structures are potential antitumor agents through different mechanisms including inhibition of several factors such as epidermal growth factor receptor (EGFR) (1,13), tubulin polymerization (14-16), poly (ADP-ribose) polymerase-1 (PARP-1) (17,18) and dihydrofolate reductase (DHFR) and thymidylate synthase (TS) (19-21) which are essential for cell growth.

In the present study 15 newly synthesized quinazolinones which were developed in our laboratory (22) were evaluated for their antibacterial, antifungal and cytotoxic activities. All of quinazolinones tested here contained aromatic substituents at position 3 and hydrazid moieties at position 2 of their structures (Fig. 1).

MATERIALS AND METHODS

The following microorganisms were used as diluted samples with broth culture: *Staphylococcus aureus*, *Bacillus subtilis*, *Listeria monocytogenes*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonella enteritidis*, *Candida albicans*, *Aspergillus niger* and *Aspergillus flavus*. All microorganisms were obtained from Persian Type Culture Collection (PTCC).

Mueller-Hinton broth (Merk, Germany) and RPMI-1640 (Gibco, Scotland) were used for growth of bacteria and fungi respectively. 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT), and dimethylsulfoxide (DMSO) were purchased from Merck, Germany. Absorbances were determined with an ELISA plate reader

(Awareness, USA). HeLa cells were purchased from Pasteure Institute, Tehran, Iran.

Antibacterial and antifungal evaluation

The *in vitro* antibacterial and antifungal activities of the synthesized compounds (Fig. 1) were studied by Microplate Alamar Blue Assay (MABA) using 96-wells microplates (23-25).

Briefly 4-5 isolated colonies were selected from an overnight bacterial culture and were diluted in broth to turbidity comparable to that of a 0.5 McFarland turbidity standard (approximately 1.0×10^8 CFU/ml). Finally inoculum density equal to 1.0×10^5 CFU/ml was




Fig. 1. Structure of new synthesized compounds studied in this work

prepared. 20 µl of each bacterial dilution was distributed in all 96 wells of microplate including positive control (containing standard antibiotic) and growth control (containing culture broth without testing materials). Then 20 µl of each concentration of the synthesized compounds were added to two neighbor wells except for positive and growth control wells. After adding alamar blue (20 µl) to all of 96 wells the total volume in each well reaches to 200 µl. The final concentrations of the tested compounds were (512, 256, 128, 64 and 32 µg/ml). After incubation, results were recorded as MIC (minimum concentration of each synthesized compound which completely inhibited growth of microorganism).

Antifungal activity was determined as antibacterial assay except with some modifications. The final size of inoculum was 1.0×10^6 CFU/ml for fungi and the turbidity was measured spectrometrically at 580 nm.

Following the MIC test, from each well which did not show growth, 50 µl of suspension was spreaded onto a plate containing appropriate medium. The plates were incubated overnight and the minimum drug concentrations able to kill 99% of bacteria and fungi were reported as Minimum Bactericidal Concentration (MBC) and Minimum Fungicidal Concentration (MFC) respectively (23). The antimicrobial tests were repeated three times.

In vitro cytotoxicity assay

HeLa cells were cultured in RPMI 1640 medium that supplemented with 50 ml heat-inactivated fetal calf serum (FCS) and sterilized by filtering through 0.22 µm filters. The HeLa cells were maintained in a humidified atmosphere of 5% CO₂, 95% air at 37°C (26,27).

The stock solutions of the synthesized compounds were prepared by dissolving in the minimum volume of dimethyl sulfoxide (DMSO) and phosphate buffered saline (PBS) was then added to reach the appropriate volume.

The cytotoxic effects of compounds against HeLa cells were evaluated using a rapid colorimetric MTT assay. Mitochondrial succinic dehydrogenase enzyme of live cells

would metabolically reduce the yellow soluble 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyl tetrazolium bromide (MTT) salt into a blue insoluble formazan product. The blue solid was dissolved in DMSO and measured spectrophotometrically in this assay. The results were compared with untreated control (26,27).

Briefly, after 2-3 subcultures, 180 µl of the cells (5×10^4 cells/ml of media) was seeded in 96 well microplates and incubated for 24 h. Each three wells of microplate was devoted to a certain concentration and 20 µl of each compound concentration was added to 180 µl of the cell suspensions in microplate wells. The final concentrations of the compounds were 1, 10 and 100 µM.

Doxorubicin was used as positive control and the wells containing DMSO (1%) and cell suspension was regarded as the negative control. The blank wells were consisted of 200 µl of the RPMI medium. The microplates were further incubated for 48 h. To evaluate cell survival, each well was then incubated with 20 µl of MTT solution (5 mg/ml in PBS) for 3 h and the media in each well was replaced with 200 µl DMSO and pipetted up and down to dissolve the formazan crystals. The absorbance of each well was measured at 540 nm using an ELISA reader. Each experiment was repeated three times. The percentage of cell viability was calculated using the following formula:

$$\% \text{ Cell Survival} = \frac{\text{Mean absorbance in drug treated wells} - \text{Mean absorbance in blank}}{\text{Mean absorbance in control wells} - \text{Mean absorbance in blank}} \times 100$$

The results are the mean of three triplicate experiments. Analysis of variance (ANOVA) followed by Tukey test was used to determine the differences between various groups. The significance level was set at $P < 0.05$.

RESULTS

Tables 1-4 present the antimicrobial effects of new 2, 3-disubstituted 4(3H)-quinazolinone derivatives. All synthesized compounds have mild to high antibacterial effects especially against Gram-negative bacteria. The most sensitive bacterium was *E. coli*. Compounds **6a** and **6b** showed weaker antibacterial effects than other compounds. All compounds showed

Table 1. The minimum inhibitory concentration (MIC) of the tested compounds against selected Gram-positive and Gram-negative bacteria

Compound	Gram-positive bacteria MIC ($\mu\text{g/ml}$)			Gram-negative bacteria MIC ($\mu\text{g/ml}$)		
	<i>S. aureus</i>	<i>L. monocytogenes</i>	<i>B. subtilis</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>Salmonella enteritidis</i>
6a	-	-	-	256	256	-
7a₁	-	-	256	64	128	128
7a₂	-	-	128	32	32	128
7a₃	-	-	512	32	32	512
7a₄	-	-	-	512	256	512
7a₅	-	-	-	128	512	-
7a₆	-	-	-	128	256	-
6b	-	-	-	256	512	-
7b₁	-	-	512	128	32	128
7b₂	-	-	512	32	32	512
7b₃	-	-	512	32	256	512
7b₄	-	-	-	32	-	-
7b₅	-	-	512	32	64	-
7b₆	-	-	-	64	512	-
7b₇	-	-	-	128	-	-

Standard antibacterial drug : Ciprofloxacin

Table 2. Minimum Bactericidal Concentration (MBC) results of the tested compounds

Compound	Gram-positive bacteria MBC ($\mu\text{g/ml}$)			Gram-negative bacteria MBC ($\mu\text{g/ml}$)		
	<i>S. aureus</i>	<i>L. monocytogenes</i>	<i>B. subtilis</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>Salmonella enteritidis</i>
7a₁	NA	NA	-	512	-	-
7a₂	NA	NA	-	512	512	-
7b₁	NA	NA	-	-	512	-
7b₃	NA	NA	-	256	-	-
7b₄	NA	NA	NA	512	NA	NA

(-): Has no bactericidal effects(NA) : Not applicable

Table 3. The minimum inhibitory concentration (MIC) of the tested compounds against selected fungi




Compounds	<i>Aspergillus flavus</i>	<i>Aspergillus niger</i>	<i>Candida albicans</i>
6a	512	512	256
7a₁	64	64	32
7a₂	32	32	32
7a₃	32	32	32
7a₄	32	64	64
7a₅	128	256	512
7a₆	512	512	512
6b	512	512	512
7b₁	64	64	64
7b₂	32	32	32
7b₃	32	32	64
7b₄	32	32	64
7b₅	64	32	32
7b₆	512	128	512
7b₇	512	512	128

Standard antifungal drug : Ketoconazole

Table 4. Minimum Fungicidal Concentration (MFC) results of the tested compounds

Compounds	<i>Aspergillus flavus</i>	<i>Aspergillus niger</i>	<i>Candida albicans</i>
7a ₁	-	-	512
7b ₂	-	-	256
7b ₃	-	-	256

(-): Has no fungicidal effects

**Fig. 2.** Cytotoxic effects of compounds **6a** and **6b** on HeLa cell line following exposure to different concentrations of compounds **6a** and **6b**, cell viability was assessed using the MTT method. Data are presented as mean \pm SD, * P <0.05, n=3, Pos-ctrl=Positive control, Neg-ctrl=Negative control**Fig. 3.** Cytotoxic effects of compounds **7a₁-7a₆** on HeLa cell line following exposure to different concentrations of compounds **7a₁-7a₆**, cell viability was assessed using the MTT method. Data are presented as mean \pm SD, * P <0.05, n=3, Pos-ctrl=Positive control, Neg-ctrl=Negative control**Fig. 4.** Cytotoxic effects of compounds **7b₁-7b₇** on HeLa cell line following exposure to different concentrations of compounds **7b₁-7b₇**, cell viability was assessed using the MTT method. Data are presented as mean \pm SD, * P <0.05, n=3, Pos-ctrl=Positive control, Neg-ctrl=Negative control**Table 5.** The percentage of cell viability following exposure to different concentrations of compounds.

Compounds	Cell survival (%)			Compound	Cell survival (%)		
	1 µM	10 µM	100 µM		1 µM	10 µM	100 µM
6a	86	73	66	6b	92	78	72
7a₁	86	74	73	7b₁	72	70	62
7a₂	90	77	68	7b₂	86	71	67
7a₃	77	75	52	7b₃	78	65	62
7a₄	74	64	58	7b₄	86	84	63
7a₅	79	74	65	7b₅	88	69	66
7a₆	87	84	65	7b₆	92	73	62
Doxorubicin	35	-	-	7b₇	90	84	80

antibacterial effects at concentrations higher than 128 µg/ml for *B. subtilis*. The tested compounds indicated no inhibitory effects against *L. monocytogenes* and *S. aureus*. Gram-positive bacteria used in the current study were insensitive to the tested compounds.

Except compounds **6a**, **7a₅**, **7a₆**, **6b**, **7b₆**, **7b₇**, all other compounds showed good antifungal effects. All three strains of fungi were sensitive to the tested compounds mostly at 32 µg/ml and there were no significant differences in the sensitivity level of the tested compounds. Synthesized compounds did not show antibacterial and antifungal effects at concentrations lower than 32 µg/ml.

Despite reasonable fungistatic and bactriostatic effects tested compounds did not show significant bactericidal and fungicidal effects according to the MBC and MFC results.

Although the synthesized compounds did not show significant cytotoxic effects, compounds **7a₃** and **7a₄** reduced cell viability to about 50% at 100 µM concentration. The cytotoxic results are shown in Figs. 2-4 and comparative results are presented in Table 5.

DISCUSSION

In comparison with some other 2,3-disubstituted quinazolinones which have some similarity in structure to the tested compounds, for example, a number of nitrogen containing groups and aromatic substituents in position 2 and 3, the studied compounds were equal or more effective in antimicrobial evaluations (28-30). It seems that the presence of nitrogen containing substituents in position 2 is beneficial for antimicrobial effects because some synthesized quinazolinones which have nitrogen containing substituents at position 2 (30) are more potent than their 3-substituted counterparts. In antimicrobial assay, compounds **6a** and **6b** (Fig. 1) are less effective than NH substituted compounds (**7a₁-7a₆** and **7b₁-7b₇**). Apparently substitution on phenyl bonded NH is necessary to have better antimicrobial effects. On the other hand it seems that the size of the substituent is important because compounds with bigger substitution showed lower antimicrobial effects. According to the results of *in vitro*

antibacterial tests which were accomplished in this study it seems that Gram-negative bacteria are more sensitive to the synthesized compounds.

Most of the studied compounds indicated high antifungal effects. The results of antifungal tests against *Candida albicans*, *Aspergillus niger* and *Aspergillus flavus* were the same and interestingly the tested compounds are more promising as potential antifungal agents.

All of the new compounds were evaluated for cytotoxic activities against the HeLa cells using MTT assay. In this assay cytotoxic activities were expressed as percentage of cell survival and were measured at 3 different concentrations. Only two compounds **7a₃**, **7a₄** reduced cell viability to about 50% at 100 µM and other compounds did not show significant cytotoxicity. Compounds **7a₃** and **7a₄** showed significant difference ($P < 0.05$) in viability in comparison to the negative control at concentrations of 1, 10, 100 µM. According to the cytotoxic results only two compounds **7a₃**, **7a₄** which have phenyl ring at position 3 and benzoyl and 4-chloro benzoyl, respectively, as N-bonded groups showed moderate cytotoxicity whereas compounds with the same groups but different substituents at position 3 did not reduce the viability of cells similar to **7a₃**, **7a₄**. Consequently it is possible to say that cell growth inhibitory effects of the compounds may be attributed to substituent property at position 3 of quinazolinone ring and the acyl group attached to the NH of phenyl hydrazine moiety.

CONCLUSION

According to the biological evaluations, tested compounds showed no significant cytotoxic effects while they possess good antifungal activities. The lack of high cytotoxicity could be beneficial as potential antifungal agents should have no toxicity on human cells. The synthesized compounds could be considered as valuable templates for further modification or derivatization to design more potent antifungal agents by the addition of different heterocyclic groups such as imidazole and triazole to the quinazolinone structure.

ACKNOWLEDGMENT

This work has been performed in the Department of Medicinal Chemistry and financially supported by the Research Council of Isfahan University of Medical Sciences.

REFERENCES

1. El-Azab AS, Al-Omar MA, Abdel-Aziz AAM, Abdel-Aziz NI, El-Sayed MAA, Aleisa AM, *et al.* Design, synthesis and biological evaluation of novel quinazoline derivatives as potential antitumor agents: Molecular docking study. *Eur J Med Chem.* 2010;45:4188-4198.
2. Ouyang G, Zhang P, Xu G, Song B, Yang S, Jin L, *et al.* Synthesis and antifungal bioactivities of 3-alkylquinazolin-4-one derivatives. *Molecules.* 2006;11:383-392.
3. Brossi A. *The alkaloids: Chemistry and Pharmacology.* New York: Academic Press; 1986. p. 131.
4. Pandeya SN, Sriram D, Nath G, De Clercq E. Synthesis, antibacterial, antifungal and anti-HIV evaluation of Schiff and Mannich bases of isatin derivatives with 3-amino-2-methylmercapto quinazolin-4 (3H)-one. *Pharm Acta Helvet.* 1999;74:11-17.
5. Panneerselvam P, Rather BA, Ravi Sankar Reddy D, Ramesh Kumar N. Synthesis and anti-microbial screening of some Schiff bases of 3-amino-6, 8-dibromo-2-phenylquinazolin-4 (3H)-ones. *Eur J Med Chem.* 2009;44:2328-2333.
6. Armarego WL. *Fused Pyrimidines.* New York: Wiley-Interscience; 1967. p. 512.
7. Aly MM, Mohamed YA, El-Bayouki KAM, Basyouni WM, Abbas SY. Synthesis of some new 4 (3H)-quinazolinone-2-carboxaldehyde thiosemicarbazones and their metal complexes and a study on their anticonvulsant, analgesic, cytotoxic and antimicrobial activities-Part-1. *Eur J Med Chem.* 2010;45:3365-3373.
8. Dinakaran M, Selvam P, DeClercq E, Sridhar SK. Synthesis, antiviral and cytotoxic activity of 6-bromo-2, 3-disubstituted-4 (3H)-quinazolinones. *Biol Pharm Bull.* 2003;26:1278-1282.
9. Raghavendra NM, Thampi P, Gurubasavarajaswamy PM, Sriram D. Synthesis and Antimicrobial Activities of Some Novel Substituted 2-Imidazolyl-N-(4-oxo-quinazolin-3 (4H)-yl)-acetamides. *Chem Pharm Bull.* 2007;55:1615-1619.
10. Hooper DC, Wolfson JS. Mode of action of the quinolone antimicrobial agents: review of recent information. *Reviews of Infectious Diseases.* 1989;11:902-911.
11. Shen L, Kohlbrenner W, Weigl D, Baranowski J. Mechanism of quinolone inhibition of DNA gyrase. Appearance of unique norfloxacin binding sites in enzyme-DNA complexes. *J Biol Chem.* 1989;264:2973.
12. Guiles J, Sun X, Critchley IA, Ochsner U, Tregay M, Stone K, *et al.* Quinazolin-2-ylamino-quinazolin-4-ols as novel non-nucleoside inhibitors of bacterial DNA polymerase III. *Bioorg Med Chem Lett.* 2009;19:800-802.
13. Lüth A, Löwe W. Syntheses of 4-(indole-3-yl)quinazolines-A new class of epidermal growth factor receptor tyrosine kinase inhibitors. *Eur J Med Chem.* 2008;43:1478-1488.
14. Raffa D, Edler MC, Daidone G, Maggio B, Merickeh M, Plescia S, *et al.* Synthesis, cytotoxicity, and inhibitory effects on tubulin polymerization of a new 3-heterocyclo substituted 2-styrylquinazolinones. *Eur J Med Chem.* 2004;39:299-304.
15. Kamal A, Bharathi EV, Reddy JS, Ramaiah MJ, Dastagiri D, Reddy MK, *et al.* Synthesis and biological evaluation of 3, 5-diaryl isoxazoline/isoxazole linked 2, 3-dihydroquinazolinone hybrids as anticancer agents. *Eur J Med Chem.* 2011;46:691-703.
16. Xia Y, Yang ZY, Hour MJ, Kuo SC, Xia P, Bastow KF, *et al.* Antitumor agents. Part 204: Synthesis and biological evaluation of substituted 2-aryl quinazolinones. *Bioorg Med Chem Lett.* 2001;11:1193-1196.
17. Iwashita A, Hattori K, Yamamoto H, Ishida J, Kido Y, Kamijo K, *et al.* Discovery of quinazolinone and quinoxaline derivatives as potent and selective poly (ADP-ribose) polymerase-1/2 inhibitors. *FEBS Lett.* 2005;579:1389-1393.
18. Orvieto F, Branca D, Giomini C, Jones P, Koch U, Ontoria JM, *et al.* Identification of substituted pyrazolo [1, 5-a] quinazolin-5 (4H)-one as potent poly (ADP-ribose) polymerase-1 (PARP-1) inhibitors. *Bioorg Med Chem Lett.* 2009;19:4196-4200.
19. Jantová S. Induction of cytotoxicity and ssDNA breaks by 9-bromo-5-morpholino-tetrazolo [1, 5-c] quinazoline in tumor cells cultured *in vitro*. *Toxicol in vitro.* 2003;17:457-463.
20. Al-Omary FAM, Abou-zeid LA, Nagi MN, Habib ESE, Abdel-Aziz AAM, El-Azab AS, *et al.* Non-classical antifolates. Part 2: Synthesis, biological evaluation, and molecular modeling study of some new 2, 6-substituted-quinazolin-4-ones. *Bioorg Med Chem.* 2010;18:2849-2863.
21. Cao SL, Feng YP, Jiang YY, Liu SY, Ding GY, Li RT, *et al.* Synthesis and *in vitro* antitumor activity of 4(3H)-quinazolinone derivatives with dithiocarbamate side chains. *Bioorg Med Chem Lett.* 2005;15:1915-1917.
22. Hassanzadeh F, Rahmani Khajouei M, Hakimelahi G, Jafari E, Khodarahmi G. Synthesis and cytotoxicity evaluation of some new 2, 3-disubstituted 4(3H)-quinazolinone derivatives. *J Res Pharm Sci.* 2012;1:23-30.
23. Schwalbe R. *Antimicrobial susceptibility testing protocols.* New York: CRC; 2007. p. 75-79.
24. Baker CN, Tenover FC. Evaluation of Alamar colorimetric broth microdilution susceptibility testing method for staphylococci and enterococci. *J Clin Microbiol.* 1996;34:2654-2659.

25. To WK, Fothergill AW, Rinaldi MG. Comparative evaluation of macrodilution and Alamar colorimetric microdilution broth methods for antifungal susceptibility testing of yeast isolates. *J Clin Microbiol.* 1995;33:2660-2664.
26. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods.* 1983;65:55-63.
27. Freshney RI. *Culture of animal cells: a manual of basic technique*: John Wiley and Sons, Inc.; 1994. p. 205-216.
28. Nanda AK, Ganguli S, Chakraborty R. Antibacterial Activity of Some 3-(Arylideneamino)-2-phenyl quinazoline-4(3H)-ones: Synthesis and Preliminary QSAR Studies. *Molecules.* 2007;12:2413-2426.
29. Kadi AA. Synthesis and antimicrobial activity of some new quinazolin-4 (3H)-one derivatives. *J Saudi Chem Soc.* 2011;15:95-100.
30. Pandey SK and Singh A. Antimicrobial studies of some novel quinazolinones fused with [1, 2, 4]-triazole, [1, 2, 4]-triazine and [1, 2, 4, 5]-tetrazine rings. *Eur J Med Chem.* 2009;44:1188-1197.